arXiv:1702.00974v1 [math.DG] 3 Feb 2017

OPTIMAL CONVERGENCE SPEED OF BERGMAN METRICS ON SYMPLECTIC MANIFOLDS

WEN LU, XIAONAN MA, AND GEORGE MARINESCU

ABSTRACT. It is known that a compact symplectic manifold endowed with a prequantum line bundle can be embedded in the projective space generated by the eigensections of low energy of the Bochner Laplacian acting on high p-tensor powers of the prequantum line bundle. We show that the Fubini-Study metrics induced by these embeddings converge at speed rate $1/p^2$ to the symplectic form.

0. INTRODUCTION

A very useful tool in the study of canonical Kähler metrics is the use of Bergman metrics to approximate arbitrary Kähler metrics in a given integral cohomology class, see e. g., [7, 11, 17].

Let (X,ω) be a compact Kähler manifold endowed with a Hermitian holomorphic line bundle (L,h^L) such that $\frac{\sqrt{-1}}{2\pi}R^L=\omega$. Since the bundle L is positive, Kodaira's theorem shows that high powers L^p give rise to holomorphic embeddings $\Phi_p : X \to \mathbb{P}(H^0(X,L^p)^*)$. The Bergman metric ω_p at level p is defined as the rescaled induced Fubini-Study metric $\frac{1}{p}\Phi_p^*\omega_{\rm FS}$, where $\omega_{\rm FS}$ is the natural Fubini-Study metric on $\mathbb{P}(H^0(X,L^p)^*)$. Tian [17] showed that ω_p converges to ω in the $\mathscr{C}^2$ topology with speed rate $p^{-1/2}$, as $p\to\infty$, that is, there exists $C>0$ such that for any $p\in\mathbb{N}^*$ we have

$$\left|\frac{1}{p}\Phi_p^*(\omega_{\rm FS})-\omega\right|_{\mathscr{C}^2}\leqslant\frac{C}{p^{1/2}}\,. \tag{0.1}$$

This was improved by Ruan [15] to convergence in $\mathscr{C}^\infty$ with speed rate p^{-1} (see also [13, Theorem 5.1.4]). Tian's result was motivated by a problem of Yau [18].

The proof of the convergence in [15, 17] is based on the diagonal expansion of the Bergman kernel up to second order. A full asymptotic expansion in powers of p in the $\mathscr{C}^\infty$ topology was obtained by Catlin [5] and Zelditch [19] as an application of Boutet de Monvel and Sjöstrand's work [4], see also [6, 14] for different approaches and generalizations. We refer to [13] for a comprehensive study of several analytic and geometric aspects of Bergman kernel. One advantage of the expansion in the $\mathscr{C}^\infty$ topology is that it easily implies the convergence of the Bergman metrics ω_p to ω with speed rate p^{-2}, see [13, (5.1.23)]. This

Date: February 6, 2017.

2010 *Mathematics Subject Classification.* 58J60, 53D50.

Key words and phrases. Symplectic Kodaira embedding, Bergman metric, Bergman kernel asymptotics, renormalized Bochner Laplacian.

W. L. supported by NNSFC No. 11401232.

X. M. partially supported by NNSFC No.11528103, ANR-14-CE25-0012-01, and funded through the Institutional Strategy of the University of Cologne within the German Excellence Initiative.

G. M. partially supported by DFG funded projects SFB TRR 191.

convergence speed is optimal and is attained only when the scalar curvature of ω is constant. Note that the scalar curvature is up to a multiplicative constant the coefficient of the second term of the Bergman kernel expansion. The purpose of this paper is to extend this optimal result to the case of symplectic manifolds.

The Bergman kernel of a holomorphic line bundle L on a complex manifold is the smooth kernel of the orthogonal projection from the space of square integrable sections on the space of holomorphic sections, or, equivalently, on the kernel of the Kodaira Laplacian $\Box^L = \overline{\partial}^L \overline{\partial}^{L*} + \overline{\partial}^{L*} \overline{\partial}^L$ on L. In order to find a suitable notion of "holomorphic section" of a prequantum line bundle on a compact *symplectic manifold*, Guillemin and Uribe [9] introduced a renormalized Bochner Laplacian $\Delta_{p,0}$ (cf. (0.5)) which reduces to $2\Box^L$ in the Kähler case.

We describe this construction in detail. Let (X, ω) be a compact symplectic manifold of real dimension $2n$. Let (L, h^L) be a Hermitian line bundle on X, and let ∇^L be a Hermitian connection on (L, h^L) with the curvature $R^L = (\nabla^L)^2$. We will assume throughout the paper that (L, h^L, ∇^L) is a prequantum line bundle of (X, ω), i.e.,

$$\frac{\sqrt{-1}}{2\pi} R^L = \omega. \tag{0.2}$$

We choose an almost complex structure J such that ω is J-invariant and $\omega(\cdot, J\cdot) > 0$. The almost complex structure J induces a splitting $TX \otimes_{\mathbb{R}} \mathbb{C} = T^{(1,0)}X \oplus T^{(0,1)}X$, where $T^{(1,0)}X$ and $T^{(0,1)}X$ are the eigenbundles of J corresponding to the eigenvalues $\sqrt{-1}$ and $-\sqrt{-1}$, respectively.

Let $g^{TX}(\cdot, \cdot) := w(\cdot, J\cdot)$ be the Riemannian metric on TX induced by ω and J. The Riemannian volume form dv_X of (X, g^{TX}) has the form $dv_X = \omega^n/n!$. The L^2-Hermitian product on the space $\mathscr{C}^\infty(X, L^p)$ of smooth sections of L^p on X, with $L^p := L^{\otimes p}$, is given by

$$\big\langle s_1, s_2 \big\rangle = \int_X \big\langle s_1, s_2 \big\rangle(x) dv_X(x). \tag{0.3}$$

Let ∇^{TX} be the Levi-Civita connection on (X, g^{TX}) with curvature R^{TX}, and let ∇^{L^p} be the connection on L^p induced by ∇^L. Let $\{e_k\}$ be a local orthonormal frame of (TX, g^{TX}). The Bochner Laplacian acting on $\mathscr{C}^\infty(X, L^p)$ is given by

$$\Delta^{L^p} = -\sum_k \Big[\big(\nabla^{L^p}_{e_k}\big)^2 - \nabla^{L^p}_{\nabla^{TX}_{e_k} e_k} \Big]. \tag{0.4}$$

Given $\Phi \in \mathscr{C}^\infty(X, \mathbb{R})$, the renormalized Bochner Laplacian is defined by

$$\Delta_{p,\Phi} = \Delta^{L^p} - 2\pi n p + \Phi. \tag{0.5}$$

By [9], [12, Corollary 1.2], there exists $C_L > 0$ independent of p such that

$$\mathrm{Spec}(\Delta_{p,\Phi}) \subset [-C_L, C_L] \cup [4\pi p - C_L, +\infty), \tag{0.6}$$

where $\mathrm{Spec}(A)$ denotes the spectrum of the operator A. Since $\Delta_{p,\Phi}$ is an elliptic operator on a compact manifold, it has discrete spectrum and its eigensections are smooth. Let $\mathcal{H}_p$ be the direct sum of eigenspaces of $\Delta_{p,\Phi}$ corresponding to the eigenvalues lying in $[-C_L, C_L]$. In mathematical physics terms, the operator $\Delta_{p,\Phi}$ is a semiclassical Schrödinger operator and the space $\mathcal{H}_p$ is the space of its bound states as $p \to \infty$. The space $\mathcal{H}_p$ proves to be an

appropriate replacement for the space of holomorphic sections $H^0(X, L^p)$ from the Kähler case. In particular, we have for p large enough (cf. [13, (8.3.3)]),

$$\dim \mathcal{H}_p = \int_X \mathrm{Td}(T^{(1,0)}X) e^{p\omega}, \tag{0.7}$$

where $\mathrm{Td}(T^{(1,0)}X)$ is the Todd class of $T^{(1,0)}X$, which corresponds to the Riemann-Roch-Hirzebruch formula from complex geometry.

Let $\mathbb{P}(\mathcal{H}_p^*)$ be the projective space associated to the dual space of $\mathcal{H}_p$; we identify $\mathbb{P}(\mathcal{H}_p^*)$ with the Grassmannian of hyperplanes in $\mathcal{H}_p$. The base locus of $\mathcal{H}_p$ is the set $\mathrm{Bl}(\mathcal{H}_p) = \big\{x \in X : s(x) = 0 \text{ for all } s \in \mathcal{H}_p\big\}$. We define the Kodaira map

$$\Phi_p : X \backslash \mathrm{Bl}(\mathcal{H}_p) \to \mathbb{P}(\mathcal{H}_p^*), \quad \Phi_p(x) = \big\{s \in \mathcal{H}_p : s(x) = 0\big\}, \tag{0.8}$$

which sends $x \in X\backslash \mathrm{Bl}(\mathcal{H})$ to the hyperplane of sections vanishing at x. Note that $\mathcal{H}_p$ is endowed with the induced L^2 Hermitian product (0.3) so there is a well-defined Fubini-Study metric g_{FS} on $\mathbb{P}(\mathcal{H}_p^*)$ with the associated form ω_{FS}.

The symplectic Kodaira embedding theorem [14, Theorem 3.6], [13, Theorem 8.3.12], states that for large p the Kodaira maps $\Phi_p : X \to \mathbb{P}(\mathcal{H}_p^*)$ are embeddings and the Bergman metrics converge to the symplectic form with speed rate p^{-1}. We note that in this case the near-diagonal expansion of the Bergman kernel is essential for the proof, in contrast to the the Kähler case, where the diagonal expansion already implies the result. Let us also observe that the results in [14, Theorem 3.6], [13, Theorem 8.3.12] are valid in a more general context, namely when g^{TX} is an arbitrary J-invariant Riemannian metric.

There exist in the literature another replacement of the notion of holomorphic section, e. g., [2, 16]. It is based on a construction by Boutet de Monvel and Guillemin [3] of a first-order pseudodifferential operator D_b on the circle bundle of L^*. The associated Szegő kernels are well defined modulo smooth operators on the associated circle bundle, even though D_b is neither canonically defined nor unique. Indeed, Boutet de Monvel–Guillemin define the Szegő kernels first, and construct the operator D_b from the Szegő kernels. Also for these spaces the Bergman metrics converge to the symplectic form with speed rate p^{-1}.

The main result of this paper is as follows.

Theorem 0.1. *Let (X, ω) be a compact symplectic manifold and (L, h^L) be a Hermitian line bundle endowed with a Hermitian connection ∇^L such that $\frac{\sqrt{-1}}{2\pi} R^L = \omega$ holds. Let J be an almost complex structure on TX such that $g^{TX}(\cdot, \cdot) := \omega(\cdot, J\cdot)$ is a J-invariant Riemannian metric on TX. Then for any $\ell \in \mathbb{N}$, there exists $C_\ell > 0$ such that*

$$\Big|\frac{1}{p}\Phi_p^*(\omega_{\mathrm{FS}}) - \omega\Big|_{\mathscr{C}^\ell} \leqslant \frac{C_\ell}{p^2}, \tag{0.9}$$

where Φ_p is the Kodaira map (0.8) *defined by the space $\mathcal{H}_p$ of bound states of the renormalized Bochner Laplacian $\Delta_{p,\Phi}$ associated with g^{TX}, ∇^L, Φ in (0.5).*

The proof is based on the near diagonal expansion of the Bergman kernel of $\mathcal{H}_p$ from [13, 14]. The sharp bound of $\mathscr{O}(p^{-2})$ is due to some remarkable cancellations of the coefficients in this expansions, reminiscent of the local properties of the curvature of Kähler metrics.

The main motivation for approximating Kähler metrics by Fubini-Study metrics comes from the questions about the existence and uniqueness of Kähler metrics of constant scalar curvature, or more generally, Kähler-Einstein metrics, see [7, 17]. It is natural to study this

questions also in the symplectic framework, for example, it is interesting to generalize to the almost-Kähler case the lower bounds on the Calabi functional given by Donaldson [8]. This is done by Lejmi and Keller [10]. Theorem 0.1 plays a crucial role in their proof in the symplectic case.

The organization of this paper is as follows. In Section 1, we recall the formal calculus on $\mathbb{C}^n$ for the model operator $\mathscr{L}$, which is the main ingredient of our approach. In Section 2, we review the asymptotic expansion of the generalized Bergman kernel. In Section 3, we reduce the proof of Theorem 0.1 to Theorem 3.3. In Section 4, we prove Theorem 3.3 and thus finish the proof of Theorem 0.1.

We shall use the following notations. For $\alpha = (\alpha_1, \ldots, \alpha_n) \in \mathbb{N}^m$, $z \in \mathbb{C}^m$, we set $|\alpha| = \sum_{j=1}^m \alpha_j$, $\alpha! = \prod_j (\alpha_j!)$ and $z^\alpha := z_1^{\alpha_1} \cdots z_m^{\alpha_m}$. Moreover, when an index variable appear twice in a single term, it means that we are summing over all its possible values.

Acknowledgments. We would like to thank Mehdi Lejmi, Julien Keller and Gábor Székelyhidi for motivating and helpful discussions which led to the writing of this paper.

1. Kernel calculus on $\mathbb{C}^n$

In this section, we recall the formal calculus on $\mathbb{C}^n$ for the model operator $\mathscr{L}$ introduced in [14, § 7.1], [13, § 4.1.6] (with $a_j = 2\pi$ therein). This calculus is the main gradient of our approach.

Let us consider the canonical coordinates $(Z_1, \ldots, Z_{2n})$ on the real vector space $\mathbb{R}^{2n}$. On the complex vector space $\mathbb{C}^n$ we consider the complex coordinates $(z_1, \ldots, z_n)$. The two sets of coordinates are linked by the relation $z_j = Z_{2j-1} + \sqrt{-1} Z_{2j}$, $j = 1, \ldots, n$.

We consider the L^2-norm

$$\| \cdot \|_{L^2} = \Big(\int_{\mathbb{R}^{2n}} |\cdot|^2 dZ \Big)^{1/2} \text{ on } \mathbb{R}^{2n}, \tag{1.1}$$

where $dZ = dZ_1 \ldots dZ_{2n}$ is the Lebesgue measure. We define the differential operators:

$$b_j = -2\frac{\partial}{\partial z_j} + \pi \overline{z}_j, \quad b_j^+ = 2\frac{\partial}{\partial \overline{z}_j} + \pi z_j, \quad b = (b_1, \ldots, b_n), \quad \mathscr{L} = \sum_{j=1}^n b_j b_j^+, \tag{1.2}$$

which extend to closed densely defined operators on $\big(L^2(\mathbb{R}^{2n}), \|\cdot\|_{L^2}\big)$. As such, b_j^+ is the adjoint of b_j and $\mathscr{L}$ defines as a densely defined self-adjoint operator on $\big(L^2(\mathbb{R}^{2n}), \|\cdot\|_{L^2}\big)$. The following result was established in [14, Theorem 1.15] (cf. also [13, Theorem 4.1.20]).

Theorem 1.1. *The spectrum of $\mathscr{L}$ on $L^2(\mathbb{R}^{2n})$ is given by*

$$\mathrm{Spec}(\mathscr{L}) = \big\{ 4\pi|\alpha| : \alpha \in \mathbb{N}^n \big\}, \tag{1.3}$$

and an orthogonal basis of the eigenspace of $4\pi|\alpha|$ is given by

$$b^\alpha \big(z^\beta \exp\big(-\pi \sum_j |z_j|^2/2 \big) \big), \quad \text{with } \beta \in \mathbb{N}^n. \tag{1.4}$$

In particular, an orthonormal basis of $\mathrm{Ker}(\mathscr{L})$ is

$$\Big\{ \phi_\beta(z) = \Big(\frac{\pi^{|\beta|}}{\beta!} \Big)^{1/2} z^\beta e^{-\pi \sum_j |z_j|^2/2} : \beta \in \mathbb{N}^n \Big\}. \tag{1.5}$$

Let $\mathscr{P}(Z,Z')$ denote the kernel of the orthogonal projection $\mathscr{P}: L^2(\mathbb{R}^{2n}) \to \mathrm{Ker}(\mathscr{L})$ with respect to dZ'. Set $\mathscr{P}^\perp = \mathrm{Id} - \mathscr{P}$.

Obviously $\mathscr{P}(Z,Z') = \sum_\beta \phi_\beta(z)\overline{\phi_\beta(z')}$, so we infer from (1.5) that

$$\mathscr{P}(Z,Z') = \exp\Big(-\frac{\pi}{2}\sum_{j=1}^{n}\big(|z_j|^2 + |z'_j|^2 - 2z_j\overline{z}'_j\big)\Big). \tag{1.6}$$

By (1.2) and (1.6), we obtain

$$\big(b_j^+\mathscr{P}\big)(Z,Z') = 0, \quad \big(b_j\mathscr{P}\big)(Z,Z') = 2\pi(\overline{z}_j - \overline{z}'_j)\mathscr{P}(Z,Z'). \tag{1.7}$$

The following commutation relations are very useful in the computations. Namely, for any polynomial $g(z,\overline{z})$ in z and $\overline{z}$, we have

$$\begin{aligned} [b_j, b_k^+] &= b_j b_k^+ - b_k^+ b_j = -4\pi\delta_{jk}, \\ [b_j, b_k] &= [b_j^+, b_k^+] = 0, \\ [g(z,\overline{z}), b_j] &= 2\frac{\partial}{\partial z_j}g(z,\overline{z}), \\ [g(z,\overline{z}), b_j^+] &= 2\frac{\partial}{\partial \overline{z}_j}g(z,\overline{z}). \end{aligned} \tag{1.8}$$

For a polynomial F in Z, Z', we denote by $F\mathscr{P}$ the operator on $L^2(\mathbb{R}^{2n})$ defined by the kernel $F(Z,Z')\mathscr{P}(Z,Z')$ and the volume form dZ.

In the calculations involving the kernel $\mathscr{P}(\cdot,\cdot)$, we prefer however to use the orthogonal decomposition of $L^2(\mathbb{R}^{2n})$ given in Theorem 1.1 and the fact that $\mathscr{P}$ is an orthogonal projection, rather than integrating against the expression (1.6) of $\mathscr{P}(\cdot,\cdot)$. This point of view helps simplify a lot the computations and understand better the operators. As an example, Theorem 1.1 implies that

$$\big(\mathscr{P}b^\alpha z^\beta \mathscr{P}\big)(Z,Z') = \begin{cases} \big(z^\beta\mathscr{P}\big)(Z,Z'), & \text{if } |\alpha| = 0, \\ 0, & \text{if } |\alpha| > 0. \end{cases} \tag{1.9}$$

We will also identify z to $\sum_j z_j\frac{\partial}{\partial z_j}$ and $\overline{z}$ to $\sum_j \overline{z}_j\frac{\partial}{\partial \overline{z}_j}$ when we consider z and $\overline{z}$ as vector fields, and

$$\mathcal{R} = \sum_j Z_j\frac{\partial}{\partial Z_j} = z + \overline{z} = Z. \tag{1.10}$$

2. Asymptotic expansion of the generalized Bergman kernel

Let a^X be the injectivity radius of (X, g^{TX}). We denote by $B^X(x,\varepsilon)$ and $B^{T_xX}(0,\varepsilon)$ the open balls in X and T_xX with center x and radius ε, respectively. Then the exponential map $T_xX \ni Z \to \exp_x^X(Z) \in X$ is a diffeomorphism from $B^{T_xX}(0,\varepsilon)$ onto $B^X(x,\varepsilon)$ for $\varepsilon \leqslant a^X$. From now on, we identify $B^{T_xX}(0,\varepsilon)$ with $B^X(x,\varepsilon)$ via the exponential map for $\varepsilon \leqslant a^X$.

We fix $x_0 \in X$. For $Z \in B^{T_{x_0}X}$ we identify (L_Z, h^L_Z) to $(L_{x_0}, h^L_{x_0})$ by parallel transport with respect to the connection ∇^L along the curve $\gamma_Z : [0,1] \ni u \to \exp_{x_0}^X(uZ)$.

In general, for functions in normal coordinates, we will add a subscript x_0 to indicate the base point $x_0 \in X$. Similarly, $P_{\mathcal{H}_p}(x,y)$ induces in terms of the above trivialization (note that

$\mathrm{End}(L^p_{x_0}) = \mathbb{C}$) a smooth function

$$\big\{(Z,Z') \in TX \times_X TX : |Z|, |Z'| < \varepsilon\big\} \ni (Z,Z') \longmapsto P_{\mathcal{H}_p,x_0}(Z,Z') \in \mathbb{C}\,,$$

which also depends smoothly on the parameter x_0.

Let us choose an orthonormal basis $\{w_j\}_{j=1}^n$ of $T^{(1,0)}_{x_0}X$. Then $e_{2j-1} = \frac{1}{\sqrt{2}}(w_j + \overline{w}_j)$ and $e_{2j} = \frac{\sqrt{-1}}{\sqrt{2}}(w_j - \overline{w}_j)$, $j = 1, \ldots, n$, forms an orthonormal basis of $T_{x_0}X$. We use coordinates on $T_{x_0}X \simeq \mathbb{R}^{2n}$ given by the identification

$$\mathbb{R}^{2n} \ni (Z_1, \ldots, Z_{2n}) \mapsto \sum_{j=1}^{2n} Z_j e_j \in T_{x_0}X. \tag{2.1}$$

In the sequel we also use complex coordinates $z = (z_1, \ldots, z_n)$ on $\mathbb{C}^n \simeq \mathbb{R}^{2n}$.

Let dv_{TX} be the Riemannian volume form on $(T_{x_0}X, g^{T_{x_0}X})$. Let $\kappa_{x_0} : T_{x_0}X \to \mathbb{R}$, $Z \mapsto \kappa_{x_0}(Z)$ be a smooth positive function defined by

$$dv_X(Z) = \kappa_{x_0}(Z) dv_{TX}(Z), \quad \kappa_{x_0}(0) = 1, \tag{2.2}$$

where the subscript x_0 of $\kappa_{x_0}(Z)$ indicates the base point $x_0 \in X$.

Rescaling $\Delta_{p,\Phi}$ and Taylor expansion. For $s \in \mathscr{C}^\infty(\mathbb{R}^{2n}, \mathbb{C})$, $Z \in \mathbb{R}^{2n}$, $|Z| \leqslant \varepsilon$, and for $t = \frac{1}{\sqrt{p}}$, set

$$\big(S_t s\big)(Z) := s(Z/t), \quad \mathscr{L}_t := S_t^{-1} \kappa^{1/2} t^2 \Delta_{p,\Phi} \kappa^{-1/2} S_t. \tag{2.3}$$

For $U \in T_{x_0}X$, we denote ∇_U the ordinary differential in direction U. Set

$$\nabla_{0,\bullet} = \nabla_\bullet + \frac{1}{2} R^L_{x_0}(Z, \cdot), \quad \mathscr{L}_0 = -\sum_{j=1}^{2n} (\nabla_{0,e_j})^2 = \sum_{j=1}^{n} b_j b_j^+ = \mathscr{L}. \tag{2.4}$$

By [14, Theorem 1.4], there exist second order differential operators $\mathcal{O}_r$ such that we have an asymptotic expansion in t when $t \to 0$,

$$\mathscr{L}_t = \mathscr{L}_0 + \sum_{r=1}^{m} t^r \mathcal{O}_r + \mathscr{O}(t^{m+1}). \tag{2.5}$$

Moreover,

$$\mathcal{O}_1(Z) = -\frac{2}{3}\big(\partial_j R^L\big)_{x_0}(\mathcal{R}, e_i) Z_j \nabla_{0,e_i} - \frac{1}{3}\big(\partial_i R^L\big)_{x_0}(\mathcal{R}, e_i), \tag{2.6}$$

and

$$\begin{aligned}
\mathcal{O}_2(Z) = {} & \frac{1}{3}\Big\langle R^{TX}_{x_0}(\mathcal{R}, e_i)\mathcal{R}, e_j\Big\rangle_{x_0} \nabla_{0,e_i}\nabla_{0,e_j} \\
& + \Big[\frac{2}{3}\Big\langle R^{TX}_{x_0}(\mathcal{R}, e_j)e_j, e_i\Big\rangle_{x_0} - \frac{1}{2}\sum_{|\alpha|=2} (\partial^\alpha R^L)_{x_0} \frac{Z^\alpha}{\alpha!}(\mathcal{R}, e_i)\Big]\nabla_{0,e_i} \\
& - \frac{1}{4}\nabla_{e_i}\Big(\sum_{|\alpha|=2} (\partial^\alpha R^L)_{x_0}(\mathcal{R}, e_i)\frac{Z^\alpha}{\alpha!}\Big) - \frac{1}{9}\sum_i \Big[\sum_j (\partial_j R^L)_{x_0}(\mathcal{R}, e_i) Z_j\Big]^2 \\
& - \frac{1}{12}\Big[\mathscr{L}_0, \Big\langle R^{TX}_{x_0}(\mathcal{R}, e_i)\mathcal{R}, e_i\Big\rangle_{x_0}\Big] + \Phi_{x_0}.
\end{aligned} \tag{2.7}$$

From (2.1) and (2.3), as in [13, Remark 4.1.8], $\mathscr{L}_t$ is a formally self-adjoint elliptic operator with respect to $\|\cdot\|_{L^2}$ on $\mathbb{R}^{2n}$ and is a smooth family of operator with respect to the parameter $x_0 \in X$. Thus $\mathscr{L}, \mathscr{L}_0$ and $\mathcal{O}_r$ in (2.5) are formally self-adjoint with respect to $\|\cdot\|_{L^2}$.

By [13, Theorem 8.3.8], the following asymptotic expansion of the generalized Bergman kernel holds.

Theorem 2.1. *There exists $J_r(Z, Z')$ polynomials in Z, Z' with the same parity as r and $\deg J_r(Z, Z') \leqslant 3r$, such that if we define*

$$\mathscr{F}_r(Z, Z') = J_r(Z, Z')\mathscr{P}(Z, Z'), \quad J_0 = 1, \tag{2.8}$$

then for any $k, \ell, m \in \mathbb{N}$, $q > 0$, there exists $C > 0$ such that if $p \geqslant 1$, $Z, Z' \in T_{x_0}X$ and $|Z|, |Z'| \leqslant \frac{q}{\sqrt{p}}$, we have

(2.9)

$$\sup_{|\alpha|+|\alpha'|\leqslant m} \Big| \frac{\partial^{|\alpha|+|\alpha'|}}{\partial Z^\alpha \partial Z'^{\alpha'}} \Big(\frac{1}{p^n} P_{\mathcal{H}_p}(Z, Z') - \sum_{r=0}^{k} \mathscr{F}_r(\sqrt{p}Z, \sqrt{p}Z') \kappa^{-\frac{1}{2}}(Z) \kappa^{-\frac{1}{2}}(Z') p^{-\frac{r}{2}} \Big) \Big|_{\mathscr{C}^\ell(X)} \leqslant C p^{-\frac{k-m+1}{2}},$$

where $\mathscr{C}^\ell(X)$ is $\mathscr{C}^\ell$-norm for the parameter $x_0 \in X$.

Moreover, by [13, (4.1.93), (8.3.45)], $\mathscr{F}_1$ and $\mathscr{F}_2$ are given by (cf. [13, (8.3.65)], [14, (1.111)])

$$\begin{aligned} \mathscr{F}_1 &= -\mathscr{P}^\perp \mathscr{L}^{-1} \mathcal{O}_1 \mathscr{P} - \mathscr{P} \mathcal{O}_1 \mathscr{L}^{-1} \mathscr{P}^\perp, \\ \mathscr{F}_2 &= \mathscr{L}^{-1} \mathscr{P}^\perp \mathcal{O}_1 \mathscr{L}^{-1} \mathscr{P}^\perp \mathcal{O}_1 \mathscr{P} - \mathscr{L}^{-1} \mathscr{P}^\perp \mathcal{O}_2 \mathscr{P} \\ &\quad + \mathscr{P} \mathcal{O}_1 \mathscr{L}^{-1} \mathscr{P}^\perp \mathcal{O}_1 \mathscr{L}^{-1} \mathscr{P}^\perp - \mathscr{P} \mathcal{O}_2 \mathscr{L}^{-1} \mathscr{P}^\perp \\ &\quad + \mathscr{P}^\perp \mathscr{L}^{-1} \mathcal{O}_1 \mathscr{P} \mathcal{O}_1 \mathscr{L}^{-1} \mathscr{P}^\perp - \mathscr{P} \mathcal{O}_1 \mathscr{L}^{-2} \mathscr{P}^\perp \mathcal{O}_1 \mathscr{P}. \end{aligned} \tag{2.10}$$

From Theorem 2.1, we get in particular [13, Theorem 8.3.3]: there exist $\boldsymbol{b}_r \in \mathscr{C}^\infty(X, \mathbb{R})$ such that for any $k, \ell \in \mathbb{N}$, there exists $C_{k,\ell} > 0$ such that

$$\Big| \frac{1}{p^n} P_{\mathcal{H}_p}(x, x) - \sum_{r=0}^{k} \boldsymbol{b}_r(x) p^{-r} \Big|_{\mathscr{C}^\ell} \leqslant C_{k,\ell}\, p^{-k-1}, \tag{2.11}$$

and

$$\boldsymbol{b}_0(x_0) = \mathscr{F}_0(0, 0) = 1, \quad \boldsymbol{b}_r(x_0) = \mathscr{F}_{2r}(0, 0), \quad \mathscr{F}_{2r+1}(0, 0) = 0. \tag{2.12}$$

3. Proof of Theorem 0.1

In this section we reduce Theorem 0.1 to Theorem 3.3. Let us fix $x_0 \in X$. As in section 2, we identify a small geodesic ball $B^X(x_0, \varepsilon)$ to $B^{T_{x_0}X}$ by means of the exponential map and we trivialize L by using a unit frame $e_L(Z)$ which is parallel with respect to ∇^L along the curve $[0, 1] \ni u \to uZ$ for $Z \in B^{T_{x_0}X}(0, \varepsilon)$.

Set $d_p := \dim \mathcal{H}_p$ and for $v = (v_1, \ldots, v_{d_p}) \in \mathbb{C}^{d_p}$, set $\|v\|^2 = \sum_{j=1}^{d_p} |v_j|^2$. We can now express the Fubini-Study metric in the homogeneous coordinate $[v] = [v_1, \ldots, v_{d_p}] \in \mathbb{P}(\mathcal{H}_p^*)$ as

$$\frac{\sqrt{-1}}{2\pi} \partial\overline{\partial} \log\big(\|v\|^2\big) = \frac{\sqrt{-1}}{2\pi} \left[\frac{1}{\|v\|^2} \sum_{j=1}^{d_p} dv_j \wedge d\overline{v}_j - \frac{1}{\|v\|^4} \sum_{j,k=1}^{d_p} \overline{v}_j v_k dv_j \wedge d\overline{v}_k \right]. \tag{3.1}$$

Let $\{s_j\}$ be an orthonormal basis of $\mathcal{H}_p$, and let $\{s^j\}$ be its dual basis. We write locally $s_j = f_j e_L^{\otimes p}$, then by (0.8), as in [13, (5.1.17)], we have

$$\Phi_p(x) = \left[\sum_{j=1}^{d_p} f_j(x) s^j\right] \in \mathbb{P}(\mathcal{H}_p^*). \tag{3.2}$$

Set

$$f^p(x,y) = \sum_{i=1}^{d_p} f_i(x)\overline{f_i}(y) \ \text{ and } \ \left|f^p(x)\right|^2 = f^p(x,x). \tag{3.3}$$

Then

$$P_{\mathcal{H}_p}(x,y) = f^p(x,y) e_L^{\otimes p}(x) \otimes e_L^{\otimes p}(y)^*, \quad \left|f^p(x)\right|^2 = P_{\mathcal{H}_p}(x,x). \tag{3.4}$$

By (3.1), (3.2) and (3.3), we get

(3.5)

$$\begin{aligned}\Phi_p^*(\omega_{FS})(x_0) &= \frac{\sqrt{-1}}{2\pi}\left[\frac{1}{|f^p|^2}\sum_{j=1}^{d_p} df_j \wedge d\overline{f_j} - \frac{1}{|f^p|^4}\sum_{j,k=1}^{d_p} \overline{f_j} f_k df_j \wedge d\overline{f_k}\right](x_0) \\ &= \frac{\sqrt{-1}}{2\pi}\Big[\left|f^p(x_0)\right|^{-2} d_x d_y f^p(x,y) - \left|f^p(x_0)\right|^{-4} d_x f^p(x,y) \wedge d_y f^p(x,y)\Big]\Big|_{x=y=x_0},\end{aligned}$$

where $\big|_{x=y=x_0}$ means the pull-back by the diagonal map $\jmath : X \to X \times X$, $x_0 \mapsto (x_0, x_0)$.

By (3.4), $P_{\mathcal{H}_p}(x,y)$ is $f^p(x,y)$ under our trivialization of L. Since we work with normal coordinates, we get from (2.2) (cf. [13, (4.1.101)])

$$\kappa(Z) = 1 + \mathscr{O}(|Z|^2). \tag{3.6}$$

By (2.9), (2.12), (3.5) and (3.6), we get

$$\begin{aligned}\frac{1}{p}\Phi_p^*(\omega_{FS})(x_0) = \frac{\sqrt{-1}}{2\pi}\Bigg\{ &\Big[\frac{1}{\mathscr{F}_0} d_x d_y \mathscr{F}_0 - \frac{1}{\mathscr{F}_0^2} d_x \mathscr{F}_0 \wedge d_y \mathscr{F}_0\Big](0,0) \\ &+ p^{-1/2}\Big[\frac{1}{\mathscr{F}_0} d_x d_y \mathscr{F}_1 - \frac{1}{\mathscr{F}_0^2}\big(d_x \mathscr{F}_1 \wedge d_y \mathscr{F}_0 + d_x \mathscr{F}_0 \wedge d_y \mathscr{F}_1\big)\Big](0,0) \\ &+ p^{-1}\Big[\frac{1}{\mathscr{F}_0} d_x d_y \mathscr{F}_2 - \frac{\mathscr{F}_2}{\mathscr{F}_0^2} d_x d_y \mathscr{F}_0 + \frac{2\mathscr{F}_2}{\mathscr{F}_0^3} d_x \mathscr{F}_0 \wedge d_y \mathscr{F}_0 \\ &\qquad - \frac{1}{\mathscr{F}_0^2}\big(d_x \mathscr{F}_0 \wedge d_y \mathscr{F}_2 + d_x \mathscr{F}_1 \wedge d_y \mathscr{F}_1 + d_x \mathscr{F}_2 \wedge d_y \mathscr{F}_0\big)\Big](0,0) \\ &+ p^{-3/2}\Big[\frac{1}{\mathscr{F}_0} d_x d_y \mathscr{F}_3 - \frac{\mathscr{F}_2}{\mathscr{F}_0^2} d_x d_y \mathscr{F}_1 - \frac{1}{\mathscr{F}_0^2}\big(d_x \mathscr{F}_0 \wedge d_y \mathscr{F}_3 \\ &\qquad + d_x \mathscr{F}_1 \wedge d_y \mathscr{F}_2 + d_x \mathscr{F}_2 \wedge d_y \mathscr{F}_1 + d_x \mathscr{F}_3 \wedge d_y \mathscr{F}_0\big) \\ &\qquad + \frac{2\mathscr{F}_2}{\mathscr{F}_0^3}\big(d_x \mathscr{F}_0 \wedge d_y \mathscr{F}_1 + d_x \mathscr{F}_1 \wedge d_y \mathscr{F}_0\big)\Big](0,0)\Bigg\} + \mathscr{O}(p^{-2}).\end{aligned} \tag{3.7}$$

From (1.6) and (2.8), we obtain

$$d_x \mathscr{F}_0(0,0) = d_y \mathscr{F}_0(0,0) = 0. \tag{3.8}$$

As J_r is a polynomial in Z, Z' with the same parity as r, we know from (1.6) and (2.8) that for $\alpha, \alpha' \in \mathbb{N}^{2n}$, there exists a polynomial $J_{r,\alpha,\alpha'}$ in Z, Z' with the same parity as $r - |\alpha| - |\alpha'|$ such that

$$\frac{\partial^{|\alpha|+|\alpha'|}}{\partial Z^{\alpha}\partial Z'^{\alpha'}}\mathscr{F}_r(Z,Z') = \big(J_{r,\alpha,\alpha'}\mathscr{P}\big)(Z,Z'). \tag{3.9}$$

In particular, (3.9) yields

$$d_x d_y \mathscr{F}_1(0,0) = d_x d_y \mathscr{F}_3(0,0) = 0, \quad d_y \mathscr{F}_2(0,0) = d_x \mathscr{F}_2(0,0) = 0. \tag{3.10}$$

By (1.6) and (2.8), we get

$$\frac{\sqrt{-1}}{2\pi}(d_x d_y \mathscr{F}_0)(0,0) = \frac{\sqrt{-1}}{2\pi}(d_x d_y \mathscr{P})(0,0) = \frac{\sqrt{-1}}{2}\sum_{j=1}^{n} dz_j \wedge d\overline{z}_j = \omega(x_0). \tag{3.11}$$

Substituting (2.12), (3.8), (3.10) and (3.11) into (3.7) yields

$$\begin{aligned}\frac{1}{p}\Phi_p^*(\omega_{FS})(x_0) = \omega(x_0) + \frac{\sqrt{-1}}{2\pi}\big(d_x d_y \mathscr{F}_2 - d_x \mathscr{F}_1 \wedge d_y \mathscr{F}_1\big)(0,0)\, p^{-1} \\ - \boldsymbol{b}_1(x_0)\omega(x_0)\, p^{-1} + \mathscr{O}(p^{-2}).\end{aligned} \tag{3.12}$$

Recall that for a tensor ψ, $\nabla^X \psi$ is the covariant derivative of ψ induced by the Levi-Civita connection ∇^{TX}. We will denote by $\langle \cdot, \cdot \rangle$ the $\mathbb{C}$-bilinear form on $TX \otimes_{\mathbb{R}} \mathbb{C}$ induced by g^{TX}.

The following observation [13, (8.3.54)] is very useful.

Lemma 3.1. *For $U \in T_{x_0}X$, $\nabla^X_U J$ is skew-adjoint and the tensor $\big\langle (\nabla^X_{\cdot} J)\cdot, \cdot \big\rangle$ is of type $\big(T^{*(1,0)}X\big)^{\otimes 3} \oplus \big(T^{*(0,1)}X\big)^{\otimes 3}$.*

Lemma 3.2. *We have*

$$(d_x \mathscr{F}_1)(0,0) = (d_y \mathscr{F}_1)(0,0) = 0. \tag{3.13}$$

Proof. By (1.2) and (2.6), we have (cf. [13, (8.3.51)])

$$\mathcal{O}_1 = -\frac{2}{3}\Big[\Big\langle (\nabla^X_{\mathcal{R}}\mathcal{J})\mathcal{R}, \frac{\partial}{\partial z_i}\Big\rangle b_i^+ - b_i\Big\langle (\nabla^X_{\mathcal{R}}\mathcal{J})\mathcal{R}, \frac{\partial}{\partial \overline{z}_i}\Big\rangle\Big], \quad \mathcal{J} = -2\pi\sqrt{-1}J. \tag{3.14}$$

From Theorem 1.1, (1.7), (3.14) and Lemma 3.1, we get (cf. [13, (8.3.67)])

$$\begin{aligned}&\Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}\mathcal{O}_1\mathscr{P}\Big)(Z,Z') \\ &\quad = -\frac{\sqrt{-1}}{3}\left\{\Big(\frac{b_i b_j}{4\pi}\Big\langle (\nabla^X_{\frac{\partial}{\partial \overline{z}_j}}J)\overline{z}', \frac{\partial}{\partial \overline{z}_i}\Big\rangle + b_i\Big\langle (\nabla^X_{\overline{z}'}J)\overline{z}', \frac{\partial}{\partial \overline{z}_i}\Big\rangle\Big)\mathscr{P}\right\}(Z,Z') \\ &\quad = -\frac{\sqrt{-1}\pi}{3}\left[\Big\langle (\nabla^X_{\overline{z}}J)\overline{z}', \overline{z}\Big\rangle + \Big\langle (\nabla^X_{\overline{z}'}J)\overline{z}', \overline{z}\Big\rangle\right]\mathscr{P}(Z,Z').\end{aligned} \tag{3.15}$$

Note that if K is an operator on $\big(\mathbb{R}^{2n}, \|\cdot\|_{L^2}\big)$ with smooth kernel $K(Z,Z')$ with respect to dZ', then the kernel $K^*(Z,Z')$ of the adjoint K^* of K, with respect to dZ', is given by

$$K^*(Z,Z') = \overline{K(Z',Z)}. \tag{3.16}$$

As $\mathscr{L}$, $\mathcal{O}_1$ are formally self-adjoint with respect to $\|\cdot\|_{L^2}$, thus $\mathscr{P}\mathcal{O}_1\mathscr{L}^{-1}\mathscr{P}^\perp$ is the adjoint of $\mathscr{L}^{-1}\mathscr{P}^\perp\mathcal{O}_1\mathscr{P}$. From Lemma 3.1, (3.15) and (3.16), we get

$$\begin{aligned}\Big(\mathscr{P}\mathcal{O}_1\mathscr{L}^{-1}\mathscr{P}^\perp\Big)(Z,Z') &= \frac{\sqrt{-1}\pi}{3}\Big[\Big\langle\big(\nabla^X_{z'}J\big)z,z'\Big\rangle+\Big\langle\big(\nabla^X_{z}J\big)z,z'\Big\rangle\Big]\mathscr{P}(Z,Z')\\ &= \frac{\sqrt{-1}\pi}{3}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_k},\frac{\partial}{\partial z_l}\Big\rangle\big(z'_jz_kz'_l+z_jz_kz'_l\big)\mathscr{P}(Z,Z').\end{aligned}\tag{3.17}$$

As the coefficient of $\mathscr{P}(Z,Z')$ in (3.15) and (3.17) are polynomials of degree 3, from (2.10), (3.15) and (3.17), we get (3.13). The proof of Lemma 3.2 is completed. □

Theorem 3.3. *The following identity holds,*

$$\frac{\sqrt{-1}}{2\pi}(d_xd_y\mathscr{F}_2)(0,0)=\boldsymbol{b}_1(x_0)\omega(x_0).\tag{3.18}$$

By Lemma 3.2, Theorem 3.3 and (3.12), we get Theorem 0.1. Section 4 is devoted to the proof of Theorem 3.3.

4. Proof of Theorem 3.3

This section is devoted to the proof of Theorem 3.3. We will compute the contribution of each term in (2.10) to $\mathscr{F}_2$. Set

$$\begin{aligned}I_1 =&\mathscr{L}^{-1}\mathscr{P}^\perp\mathcal{O}_1\mathscr{L}^{-1}\mathscr{P}^\perp\mathcal{O}_1\mathscr{P},\quad I_2=-\mathscr{L}^{-1}\mathscr{P}^\perp\mathcal{O}_2\mathscr{P},\\ I_3 =&\mathscr{P}\mathcal{O}_1\mathscr{L}^{-1}\mathscr{P}^\perp\mathcal{O}_1\mathscr{L}^{-1}\mathscr{P}^\perp,\quad I_4=-\mathscr{P}\mathcal{O}_2\mathscr{L}^{-1}\mathscr{P}^\perp,\\ I_5 =&\mathscr{P}^\perp\mathscr{L}^{-1}\mathcal{O}_1\mathscr{P}\mathcal{O}_1\mathscr{L}^{-1}\mathscr{P}^\perp,\quad I_6=-\mathscr{P}\mathcal{O}_1\mathscr{L}^{-2}\mathscr{P}^\perp\mathcal{O}_1\mathscr{P}.\end{aligned}\tag{4.1}$$

For $j\in\{1,\dots,6\}$, let $I_j(Z,Z')$ be the smooth kernel of the operator I_j with respect to dZ'. By (2.10),

$$(d_xd_y\mathscr{F}_2)(0,0)=\sum_{j=1}^{6}(d_xd_yI_j)(0,0).\tag{4.2}$$

In the context of (3.16), if we denote $b_{jk}=\dfrac{\partial^2K}{\partial Z_j\partial Z'_k}(Z,Z')\big|_{Z=Z'=0}$, then

$$\begin{aligned}\big(d_Zd_{Z'}K^*\big)(0,0)&=\sum_{j,k}dZ_j\wedge dZ_k\,\frac{\partial^2K^*}{\partial Z_j\partial Z'_k}(Z,Z')\big|_{Z=Z'=0}\\ &=\sum_{j<k}\big(\overline{b}_{kj}-\overline{b}_{jk}\big)dZ_j\wedge dZ_k=-\overline{\big(d_Zd_{Z'}K\big)(0,0)}.\end{aligned}\tag{4.3}$$

As $\mathcal{O}_r$ in (2.5) are formally self-adjoint with respect to $\|\cdot\|_{L^2}$, we get by (4.1) that I_1 and I_2 are adjoints of I_3 and I_4, respectively, as operators acting on $\big(\mathbb{R}^{2n},\|\cdot\|_{L^2}\big)$. Hence by (4.3),

$$(d_xd_yI_3)(0,0)=-\overline{(d_xd_yI_1)(0,0)},\quad (d_xd_yI_4)(0,0)=-\overline{(d_xd_yI_2)(0,0)}.\tag{4.4}$$

4.1. **Evaluation of** $(d_xd_yI_j)(0,0)$ **for** $j=1,3,5,6$**.** To simplify the notation, for polynomials Q_1, Q_2 in Z, Z', we will denote

$$(Q_1\mathscr{P})(Z,Z')\sim(Q_2\mathscr{P})(Z,Z'), \tag{4.5}$$

if the constant coefficient and for all j the coefficient of Z'_j in Q_1-Q_2 as a polynomial in Z' are zero; we denote

$$(Q_1\mathscr{P})(Z,Z')\approx(Q_2\mathscr{P})(Z,Z'), \tag{4.6}$$

if (4.5) holds and the constant coefficient and for all j the coefficient of Z_j in Q_1-Q_2 as a polynomial in Z are zero.

Set

$$\mathscr{J}_{jir}:=\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_i},\frac{\partial}{\partial z_r}\Big\rangle,\quad \mathscr{J}_{\overline{j}\,\overline{ir}}:=\Big\langle\big(\nabla^X_{\frac{\partial}{\partial \overline{z}_j}}J\big)\frac{\partial}{\partial \overline{z}_i},\frac{\partial}{\partial \overline{z}_r}\Big\rangle=\overline{\mathscr{J}}_{jir}. \tag{4.7}$$

From (1.7), (3.14), Lemma 3.1 and (3.15), we get

$$\begin{aligned}
&\Big(\mathcal{O}_1\mathscr{L}^{-1}\mathscr{P}^\perp\mathcal{O}_1\mathscr{P}\Big)(Z,Z')\\
&=\frac{4\pi}{9}\bigg\{\Big(\Big\langle\big(\nabla^X_z J\big)z,\frac{\partial}{\partial z_i}\Big\rangle b_i^+-b_i\Big\langle\big(\nabla^X_{\overline{z}}J\big)\overline{z},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\Big)\\
&\qquad\times\Big(\frac{b_jb_k}{4\pi}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_k}}J\big)\overline{z}',\frac{\partial}{\partial\overline{z}_j}\Big\rangle+b_j\Big\langle\big(\nabla^X_{\overline{z}'}J\big)\overline{z}',\frac{\partial}{\partial\overline{z}_j}\Big\rangle\Big)\mathscr{P}\bigg\}(Z,Z')\\
&\sim\frac{1}{9}\bigg\{\Big[\Big\langle\big(\nabla^X_z J\big)z,\frac{\partial}{\partial z_i}\Big\rangle b_i^+-b_i\Big\langle\big(\nabla^X_{\overline{z}}J\big)\overline{z},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\Big]b_jb_k\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_k}}J\big)\overline{z}',\frac{\partial}{\partial\overline{z}_j}\Big\rangle\mathscr{P}\bigg\}(Z,Z')\\
&\sim\frac{1}{9}\bigg\{\Big\langle\big(\nabla^X_z J\big)z,\frac{\partial}{\partial z_i}\Big\rangle\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_k}}J\big)\overline{z}',\frac{\partial}{\partial\overline{z}_j}\Big\rangle b_i^+b_jb_k\mathscr{P}\bigg\}(Z,Z'),
\end{aligned} \tag{4.8}$$

in the last equation of (4.8), we use $b_i\Big\langle\big(\nabla^X_{\overline{z}}J\big)\overline{z},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\mathscr{P}=\Big\langle\big(\nabla^X_{\overline{z}}J\big)\overline{z},-2\pi\overline{z}'\Big\rangle\mathscr{P}$.

By Theorem 1.1, (1.7) and (1.8), we get $b_i^+b_jb_k=b_jb_kb_i^++4\pi\big(\delta_{ij}b_k+\delta_{ik}b_j\big)$ and

$$\begin{aligned}
\mathscr{L}^{-1}\mathscr{P}^\perp\big(z_sz_tb_i^+b_jb_k\mathscr{P}\big)&=4\pi\mathscr{L}^{-1}\mathscr{P}^\perp\big(z_sz_t(\delta_{ij}b_k+\delta_{ik}b_j)\mathscr{P}\big)\\
&=4\pi\mathscr{L}^{-1}\mathscr{P}^\perp\big((\delta_{ij}b_k+\delta_{ik}b_j)z_sz_t\mathscr{P}\big)\\
&=(\delta_{ij}b_k+\delta_{ik}b_j)z_sz_t\mathscr{P}.
\end{aligned} \tag{4.9}$$

By (1.7), (1.8), (4.1), (4.8) and (4.9), we get

$$\begin{aligned}
I_1(Z,Z')&\sim\frac{1}{9}\mathscr{J}_{sti}\mathscr{J}_{\overline{k}\,\overline{lj}}\big((\delta_{ij}b_kz_sz_t+\delta_{ik}b_jz_sz_t)\overline{z}'_l\mathscr{P}\big)(Z,Z')\\
&=\frac{1}{9}\mathscr{J}_{sti}\mathscr{J}_{\overline{k}\,\overline{lj}}\Big[-2\delta_{ij}\delta_{ks}z_t-2\delta_{ij}\delta_{kt}z_s+2\pi\delta_{ij}z_sz_t(\overline{z}_k-\overline{z}'_k)\\
&\qquad-2\delta_{ik}\delta_{js}z_t-2\delta_{ik}\delta_{jt}z_s+2\pi\delta_{ik}z_sz_t(\overline{z}_j-\overline{z}'_j)\Big]\overline{z}'_l\mathscr{P}(Z,Z').
\end{aligned} \tag{4.10}$$

Recall that $\mathscr{J}_{sti}$ is anti-symmetric on t and i, thus the contribution of $-2\delta_{ij}\delta_{kt}z_s-2\delta_{ik}\delta_{jt}z_s$ in (4.10) is zero. Thus from (4.10), we get

$$I_1(Z,Z')\approx-\frac{2}{9}\mathscr{J}_{sri}\mathscr{J}_{\overline{k}\,\overline{qj}}\big(\delta_{ik}\delta_{js}+\delta_{ij}\delta_{ks}\big)z_r\overline{z}'_q\mathscr{P}(Z,Z'). \tag{4.11}$$

By (1.6), Lemma 3.1 and (4.11), we get

$$(d_xd_yI_1)(0,0)=-\frac{2}{9}\mathscr{J}_{jir}\big(\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}+\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}\big)dz_r\wedge d\overline{z}_q. \tag{4.12}$$

From (4.4), (4.7) and (4.12), we get

$$(d_xd_yI_3)(0,0)=(d_xd_yI_1)(0,0). \tag{4.13}$$

By (1.6), (3.15), (3.17) and (4.1), we get

(4.14)

$$\begin{aligned}I_5(Z,Z')&\sim\frac{\pi^2}{9}\bigg\{\Big(\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}}}J\big)\overline{z}''+\big(\nabla^X_{\frac{\partial}{\partial\overline{z}''}}J\big)\overline{z}'',\overline{z}\Big\rangle\mathscr{P}\Big)\circ\Big(\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_k},\frac{\partial}{\partial z_l}\Big\rangle z_j''z_k''z_l'\mathscr{P}\Big)\bigg\}(Z,Z')\\&\approx\frac{\pi^2}{9}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_s}}J\big)\frac{\partial}{\partial\overline{z}_t},\overline{z}\Big\rangle\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_k},z'\Big\rangle\Big\{\mathscr{P}\circ\big(\overline{z}_s''\overline{z}_t''z_j''z_k''\mathscr{P}\big)\Big\}(Z,Z')\\&\approx\frac{\pi^2}{9}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_s}}J\big)\frac{\partial}{\partial\overline{z}_t},\overline{z}\Big\rangle\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_k},z'\Big\rangle\mathscr{P}(Z,Z')\Big(\mathscr{P}\circ\big(\overline{z}_s\overline{z}_tz_jz_k\mathscr{P}\big)\Big)(0,0),\end{aligned}$$

where in the last equation we use $\mathscr{P}(0,0)=1$, since we need to compute the constant coefficient of $\mathscr{P}$ in $\mathscr{P}\circ\big(\overline{z}_s''\overline{z}_t''z_j''z_k''\mathscr{P}\big)$.

By (1.7) and (1.8), we get

$$\begin{aligned}(\overline{z}_s\overline{z}_tz_jz_k\mathscr{P})(Z,0)&=\frac{1}{4\pi^2}(z_jz_kb_sb_t\mathscr{P})(Z,0),\\z_jz_kb_sb_t&=b_sb_tz_jz_k+2\delta_{js}b_tz_k+2\delta_{jt}b_sz_k\\&\quad+2\delta_{ks}b_tz_j+2\delta_{kt}b_sz_j+4\delta_{jt}\delta_{ks}+4\delta_{js}\delta_{kt}.\end{aligned} \tag{4.15}$$

From Theorem 1.1 and (4.15), we get

$$\begin{aligned}\Big(\mathscr{P}\circ\big(\overline{z}_s\overline{z}_tz_jz_k\mathscr{P}\big)\Big)(0,0)&=\frac{1}{4\pi^2}(4\delta_{jt}\delta_{ks}+4\delta_{js}\delta_{kt})\mathscr{P}(0,0)\\&=\frac{1}{\pi^2}(\delta_{jt}\delta_{ks}+\delta_{js}\delta_{kt}).\end{aligned} \tag{4.16}$$

From (4.7), (4.14) and (4.16), we obtain

$$(d_xd_yI_5)(0,0)=-\frac{1}{9}\mathscr{J}_{jir}\big(\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}+\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}\big)dz_r\wedge d\overline{z}_q. \tag{4.17}$$

By (3.15), (3.17) and (4.1), we get

$$\begin{aligned}I_6(Z,Z')&\approx-\frac{\pi^2}{9}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)z,\frac{\partial}{\partial z_l}\Big\rangle\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_s}}J\big)\overline{z}',\frac{\partial}{\partial\overline{z}_t}\Big\rangle\Big(\mathscr{P}\circ\big(z_j''z_l''\overline{z}_s''\overline{z}_t''\mathscr{P}\big)\Big)(Z,Z')\\&\approx-\frac{\pi^2}{9}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)z,\frac{\partial}{\partial z_k}\Big\rangle\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_s}}J\big)\overline{z}',\frac{\partial}{\partial\overline{z}_t}\Big\rangle\mathscr{P}(Z,Z')\Big(\mathscr{P}\circ\big(z_jz_k\overline{z}_s\overline{z}_t\mathscr{P}\big)\Big)(0,0).\end{aligned} \tag{4.18}$$

Thus by (4.7), (4.16) and (4.18), we get

$$\begin{aligned}(d_xd_yI_6)(0,0)&=-\frac{1}{9}\mathscr{J}_{jrk}\mathscr{J}_{\overline{s}\,\overline{q}\overline{t}}\big(\delta_{jt}\delta_{ks}+\delta_{js}\delta_{kt}\big)dz_r\wedge d\overline{z}_q\\&=-\frac{1}{9}\mathscr{J}_{jir}\big(\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}+\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}\big)dz_r\wedge d\overline{z}_q.\end{aligned} \tag{4.19}$$

4.2. **Evaluation of** $(d_xd_yI_2)(0,0)$**: part I.** Recall that [14, Lemma 2.1],

$$\begin{aligned}(4.20)\quad \mathcal{O}_2\mathscr{P} = \Big\{&\frac{1}{3}b_ib_j\Big\langle R^{TX}_{x_0}(\mathcal{R},\frac{\partial}{\partial\overline{z}_i})\mathcal{R},\frac{\partial}{\partial\overline{z}_j}\Big\rangle + \frac{1}{2}b_i\Big[\sum_{|\alpha|=2}\big(\partial^\alpha R^L\big)(\mathcal{R},\frac{\partial}{\partial\overline{z}_i})\frac{Z^\alpha}{\alpha!}\Big]\\ &+\frac{4}{3}b_j\Big[\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big)\mathcal{R},\frac{\partial}{\partial\overline{z}_j}\Big\rangle - \Big\langle R^{TX}\Big(\mathcal{R},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\Big]\\ &-2\pi\sqrt{-1}\Big\langle\big(\nabla^X\nabla^XJ\big)_{(\mathcal{R},\mathcal{R})}\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big\rangle + 4\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial z_j}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\Big\}\mathscr{P}\\ &+\Big[-\frac{1}{3}\mathscr{L}\Big\langle R^{TX}\Big(\mathcal{R},\frac{\partial}{\partial z_j}\Big)\mathcal{R},\frac{\partial}{\partial\overline{z}_j}\Big\rangle + \frac{4\pi^2}{9}\big|(\nabla^X_{\mathcal{R}}J)\mathcal{R}\big|^2+\Phi_{x_0}\Big]\mathscr{P}.\end{aligned}$$

Set

$$\begin{aligned}(4.21)\quad I_{21}(Z,Z') &= \frac{1}{3}\Big(\mathscr{L}^{-1}\mathscr{P}^\perp b_ib_j\Big\langle R^{TX}_{x_0}(\mathcal{R},\frac{\partial}{\partial\overline{z}_i})\mathcal{R},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\mathscr{P}\Big)(Z,Z'),\\ I_{22}(Z,Z') &= \frac{1}{2}\Big(\mathscr{L}^{-1}\mathscr{P}^\perp b_i\Big[\sum_{|\alpha|=2}\big(\partial^\alpha R^L\big)\frac{Z^\alpha}{\alpha!}(\mathcal{R},\frac{\partial}{\partial\overline{z}_i})\Big]\mathscr{P}\Big)(Z,Z'),\\ I_{23}(Z,Z') &= \frac{4}{3}\Big\{\mathscr{L}^{-1}b_j\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big)\mathcal{R} - R^{TX}\Big(\mathcal{R},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\mathscr{P}\Big\}(Z,Z'),\\ I_{24}(Z,Z') &= -2\pi\sqrt{-1}\Big(\mathscr{L}^{-1}\mathscr{P}^\perp\Big\langle\big(\nabla^X\nabla^XJ\big)_{(\mathcal{R},\mathcal{R})}\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\mathscr{P}\Big)(Z,Z'),\\ I_{25}(Z,Z') &= -\frac{1}{3}\Big(\mathscr{P}^\perp\mathscr{L}^{-1}\mathscr{L}\Big\langle R^{TX}\Big(\mathcal{R},\frac{\partial}{\partial z_i}\Big)\mathcal{R},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\mathscr{P}\Big)(Z,Z'),\\ I_{26}(Z,Z') &= \frac{4\pi^2}{9}\Big(\mathscr{L}^{-1}\mathscr{P}^\perp\big|(\nabla^X_{\mathcal{R}}J)\mathcal{R}\big|^2\mathscr{P}\Big)(Z,Z').\end{aligned}$$

By Theorem 1.1, $\mathscr{P}^\perp\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial z_j}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\mathscr{P} = \mathscr{P}^\perp\Phi_{x_0}\mathscr{P} = 0$. Thus by (4.1), (4.20) and (4.21), we get

$$(4.22)\qquad -I_2(Z,Z') = \sum_{j=1}^{6} I_{2j}(Z,Z').$$

We evaluate first the contribution of I_{2j}, $j=1,3,5,6$, in $(d_xd_yI_2)(0,0)$. We recall the following well-known symmetry properties of the curvature R^{TX}: for $U,V,W,Y\in TX$, we have

$$\begin{aligned}(4.23)\quad &\Big\langle R^{TX}(U,V)W,Y\Big\rangle = \Big\langle R^{TX}(W,Y)U,V\Big\rangle,\\ &R^{TX}(U,V)W + R^{TX}(V,W)U + R^{TX}(W,U)V = 0.\end{aligned}$$

Using (1.8) and (4.23), we have

$$\begin{aligned}(4.24)\quad b_ib_j\Big\langle R^{TX}\Big(\mathcal{R},\frac{\partial}{\partial\overline{z}_i}\Big)\mathcal{R},\frac{\partial}{\partial\overline{z}_j}\Big\rangle &= b_ib_j\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_s},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_t},\frac{\partial}{\partial\overline{z}_j}\Big\rangle z_sz_t\\ &+2b_ib_j\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_s},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial\overline{z}_t},\frac{\partial}{\partial\overline{z}_j}\Big\rangle z_s\overline{z}_t + b_ib_j\Big\langle R^{TX}\Big(\frac{\partial}{\partial\overline{z}_s},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial\overline{z}_t},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\overline{z}_s\overline{z}_t.\end{aligned}$$

By (1.7) and (1.8), we get

$$z_s\overline{z}_t\mathscr{P}(Z,Z')=z_s(\frac{b_t}{2\pi}+\overline{z}'_t)\mathscr{P}(Z,Z')=\Big(\frac{b_t}{2\pi}z_s+\frac{\delta_{st}}{\pi}+z_s\overline{z}'_t\Big)\mathscr{P}(Z,Z'). \tag{4.25}$$

By Theorem 1.1, (4.21), (4.24) and (4.25), we get

$$\begin{aligned}3I_{21}(Z,Z')=&\frac{1}{8\pi}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_s},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_t},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\Big(b_ib_jz_sz_t\mathscr{P}\Big)(Z,Z')\\&+\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_s},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial\overline{z}_t},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\Big[\big(\frac{b_ib_jb_t}{12\pi^2}z_s+\frac{b_ib_j}{4\pi}z_s\overline{z}'_t\big)\mathscr{P}\Big](Z,Z')\\&+\frac{1}{4\pi^2}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_k},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial\overline{z}_k},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\Big(b_ib_j\mathscr{P}\Big)(Z,Z')+I_{27}(Z,Z'),\end{aligned} \tag{4.26}$$

where

$$I_{27}(Z,Z')=\Big\langle R^{TX}\Big(\frac{\partial}{\partial\overline{z}_s},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial\overline{z}_t},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}b_ib_j\overline{z}_s\overline{z}_t\mathscr{P}\Big)(Z,Z'). \tag{4.27}$$

Note that by Theorem 1.1, (1.7) and (1.8),

$$\begin{aligned}4\pi^2\mathscr{L}^{-1}\mathscr{P}^{\perp}b_ib_j\overline{z}_s\overline{z}_t\mathscr{P}&=\mathscr{L}^{-1}\mathscr{P}^{\perp}b_ib_j\big(b_s+2\pi\overline{z}'_s\big)\big(b_t+2\pi\overline{z}'_t\big)\mathscr{P}\\&=\Big[\frac{1}{16\pi}b_ib_jb_sb_t+\frac{1}{6}b_ib_j\big(b_s\overline{z}'_t+b_t\overline{z}'_s\big)+\frac{\pi}{2}b_ib_j\overline{z}'_s\overline{z}'_t\Big]\mathscr{P}.\end{aligned} \tag{4.28}$$

Thus, from (1.7), (4.6) and (4.28), we get

$$I_{27}(Z,Z')\approx 0. \tag{4.29}$$

From (1.7) and (1.8), we get

$$\begin{aligned}\big(b_ib_j\mathscr{P}\big)(Z,Z')&=4\pi^2\big(\overline{z}_i-\overline{z}'_i\big)\big(\overline{z}_j-\overline{z}'_j\big)\mathscr{P}(Z,Z'),\\\big(b_ib_jz_s\overline{z}'_t\mathscr{P}\big)(Z,Z')&=\Big[-4\pi\delta_{js}\overline{z}'_t\big(\overline{z}_i-\overline{z}'_i\big)-4\pi\delta_{is}\overline{z}'_t(\overline{z}_j-\overline{z}'_j)\\&\qquad+4\pi^2z_s\overline{z}'_t\big(\overline{z}_i-\overline{z}'_i\big)\big(\overline{z}_j-\overline{z}'_j\big)\Big]\mathscr{P}(Z,Z'),\\\big(b_ib_jz_sz_t\mathscr{P}\big)(Z,Z')&=\Big[4\delta_{it}\delta_{js}-4\pi\delta_{js}z_t\big(\overline{z}_i-\overline{z}'_i\big)+4\delta_{jt}\delta_{is}-4\pi\delta_{jt}z_s\big(\overline{z}_i-\overline{z}'_i\big)\\&\qquad-4\pi\delta_{is}z_t\big(\overline{z}_j-\overline{z}'_j\big)-4\pi\delta_{it}z_s\big(\overline{z}_j-\overline{z}'_j\big)\\&\qquad+4\pi^2z_sz_t\big(\overline{z}_j-\overline{z}'_j\big)\big(\overline{z}_i-\overline{z}'_i\big)\Big]\mathscr{P}(Z,Z'),\end{aligned} \tag{4.30}$$

and

$$\Big(b_ib_jb_tz_s\mathscr{P}\Big)(Z,Z')=\Big[\big(-2\delta_{ts}b_ib_j-2\delta_{js}b_ib_t-2\delta_{is}b_jb_t+z_sb_ib_jb_t\big)\mathscr{P}\Big](Z,Z'). \tag{4.31}$$

By (1.7), (4.30) and (4.31), we get

$$\big(d_xd_y(b_ib_j\mathscr{P})\big)(0,0)=0,\quad\big(d_xd_y(b_ib_jb_tz_s\mathscr{P})\big)(0,0)=0. \tag{4.32}$$

Substituting (3.11), (4.23), (4.29)–(4.32) into (4.26), we obtain

$$\begin{aligned}(d_xd_yI_{21})(0,0)=&-\frac{\sqrt{-1}}{3}\Big\langle 2R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_i}-R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\omega(x_0)\\&+\frac{1}{3}\Big\langle 2R^{TX}\Big(\frac{\partial}{\partial z_r},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial z_j}+R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_r}\Big)\frac{\partial}{\partial\overline{z}_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle dz_r\wedge d\overline{z}_q\\&+\frac{1}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_r}\Big)\frac{\partial}{\partial\overline{z}_j}+R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial\overline{z}_r},\frac{\partial}{\partial\overline{z}_q}\Big\rangle d\overline{z}_r\wedge d\overline{z}_q.\end{aligned}\tag{4.33}$$

By (4.21),

$$\begin{aligned}\frac{3}{4}I_{23}=&\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_s}-R^{TX}\Big(\frac{\partial}{\partial z_s},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\mathscr{L}^{-1}b_jz_s\mathscr{P}\\&+\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial\overline{z}_s}-R^{TX}\Big(\frac{\partial}{\partial\overline{z}_s},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\mathscr{L}^{-1}b_j\overline{z}_s\mathscr{P}.\end{aligned}\tag{4.34}$$

By Theorem 1.1, (1.7) and (1.8),

$$\begin{aligned}\Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}b_jz_s\mathscr{P}\Big)(Z,Z')&=\frac{1}{4\pi}\Big(b_jz_s\mathscr{P}\Big)(Z,Z')\\&=\frac{1}{4\pi}\Big(-2\delta_{js}+2\pi z_s\big(\overline{z}_j-\overline{z}'_j\big)\Big)\mathscr{P}(Z,Z').\end{aligned}\tag{4.35}$$

Note that by (1.7), $\overline{z}_s\mathscr{P}=\Big(\frac{b_s}{2\pi}+\overline{z}'_s\Big)\mathscr{P}$. Thus from Theorem 1.1, we get

$$\begin{aligned}\Big(\mathscr{L}^{-1}b_j\overline{z}_s\mathscr{P}\Big)(Z,Z')&=\Big[\big(\frac{b_jb_s}{16\pi^2}+\frac{b_j}{4\pi}\overline{z}'_s\big)\mathscr{P}\Big](Z,Z')\\&=\Big[\frac{1}{4}\big(\overline{z}_j-\overline{z}'_j\big)\big(\overline{z}_s-\overline{z}'_s\big)+\frac{1}{2}\overline{z}'_s\big(\overline{z}_j-\overline{z}'_j\big)\Big]\mathscr{P}(Z,Z').\end{aligned}\tag{4.36}$$

As in (4.32), we get

$$\big(d_xd_y(\mathscr{L}^{-1}b_j\overline{z}_s\mathscr{P})\big)(0,0)=\frac{1}{2}d\overline{z}_j\wedge d\overline{z}_s.\tag{4.37}$$

From (3.11), (4.23), (4.34), (4.35) and (4.37), we get

$$\begin{aligned}(d_xd_yI_{23})(0,0)=&\frac{4\sqrt{-1}}{3}\Big\langle R^{TX}(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i})\frac{\partial}{\partial z_i}-2R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\omega(x_0)\\&-\frac{2}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_r},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_i}+2R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial z_r}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_q}\Big\rangle dz_r\wedge d\overline{z}_q\\&-\frac{2}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial\overline{z}_r}+R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_r}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_q}\Big\rangle d\overline{z}_r\wedge d\overline{z}_q.\end{aligned}\tag{4.38}$$

Clearly, by Theorem 1.1 and (4.21),

$$\begin{aligned}-3I_{25}(Z,Z')=&\Big(\mathscr{P}^{\perp}\Big\langle R^{TX}\Big(\mathcal{R},\frac{\partial}{\partial z_i}\Big)\mathcal{R},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\mathscr{P}\Big)(Z,Z')\\=&\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial\overline{z}_k}+R^{TX}\Big(\frac{\partial}{\partial\overline{z}_k},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\Big(\mathscr{P}^{\perp}\circ\big(z_j\overline{z}_k\mathscr{P}\big)\Big)(Z,Z')\\&+\Big\langle R^{TX}\Big(\frac{\partial}{\partial\overline{z}_j},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial\overline{z}_k},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\Big(\mathscr{P}^{\perp}\circ\big(\overline{z}_j\overline{z}_k\mathscr{P}\big)\Big)(Z,Z').\end{aligned}\tag{4.39}$$

From Theorem 1.1, (1.7) and (4.25), we get

$$\Big(\mathscr{P}^{\perp}\circ\big(z_j\overline{z}_k\mathscr{P}\big)\Big)(Z,Z')=\frac{1}{2\pi}\big(b_kz_j\mathscr{P}\big)(Z,Z')=\Big(-\frac{1}{\pi}\delta_{jk}+z_j(\overline{z}_k-\overline{z}'_k)\Big)\mathscr{P}(Z,Z'), \tag{4.40}$$

and

$$\begin{aligned}\Big(\mathscr{P}^{\perp}\circ\big(\overline{z}_j\overline{z}_k\mathscr{P}\big)\Big)(Z,Z')&=\Big\{\mathscr{P}^{\perp}\Big[\frac{1}{4\pi^2}b_jb_k+\frac{1}{2\pi}(b_j\overline{z}'_k+b_k\overline{z}'_j)\Big]\mathscr{P}\Big\}(Z,Z')\\&=\Big((\overline{z}_j-\overline{z}'_j)(\overline{z}_k-\overline{z}'_k)+\overline{z}'_k(\overline{z}_j-\overline{z}'_j)+\overline{z}'_j(\overline{z}_k-\overline{z}'_k)\Big)\mathscr{P}(Z,Z')\\&=\big(\overline{z}_j\overline{z}_k-\overline{z}'_j\overline{z}'_k\big)\mathscr{P}(Z,Z').\end{aligned} \tag{4.41}$$

As in [13, (8.3.56), (8.3.63)], we have

$$\begin{aligned}&\big|\nabla^XJ\big|^2=\sum_{i,j}\big|(\nabla^X_{e_i}J)e_j\big|^2=8\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_i}}J\big)\frac{\partial}{\partial z_j},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\frac{\partial}{\partial\overline{z}_j}\Big\rangle,\\&\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial z_j}\Big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle=\frac{1}{32}\big|\nabla^XJ\big|^2.\end{aligned} \tag{4.42}$$

By (3.11), (4.23), (4.39), (4.40), (4.41) and (4.42), we get

$$\begin{aligned}(d_xd_yI_{25})(0,0)&=-\frac{1}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial\overline{z}_k}+R^{TX}\Big(\frac{\partial}{\partial\overline{z}_k},\frac{\partial}{\partial z_i}\Big)\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\\&\qquad\times\Big(2\sqrt{-1}\delta_{jk}\,\omega(x_0)-dz_j\wedge d\overline{z}_k\Big)\\&=\Big[-\frac{1}{96}\big|\nabla^XJ\big|^2+\frac{1}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\Big]2\sqrt{-1}\omega(x_0)\\&\qquad+\frac{1}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial z_r}\Big)\frac{\partial}{\partial\overline{z}_i}-R^{TX}\Big(\frac{\partial}{\partial z_r},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_q}\Big\rangle dz_r\wedge d\overline{z}_q.\end{aligned} \tag{4.43}$$

Then by Lemma 3.1 and (4.21), we get

$$9I_{26}(Z,Z')=8\pi^2\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_i}}J\big)\frac{\partial}{\partial z_j},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_s}}J\big)\frac{\partial}{\partial\overline{z}_t}\Big\rangle\Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}z_iz_j\overline{z}_s\overline{z}_t\mathscr{P}\Big)(Z,Z'). \tag{4.44}$$

By Theorem 1.1, (1.7), (1.8) and (4.15), we get

$$\Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}z_iz_jb_s\mathscr{P}\Big)(Z,Z')=\frac{b_s}{4\pi}z_iz_j\mathscr{P}(Z,Z')=\frac{1}{4\pi}\big(-2\delta_{is}z_j-2\delta_{js}z_i+z_iz_jb_s\big)\mathscr{P}(Z,Z'), \tag{4.45}$$

$$\begin{aligned}&\Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}z_iz_jb_sb_t\mathscr{P}\Big)(Z,Z')=\frac{1}{2\pi}\Big(\frac{b_sb_t}{4}z_iz_j+\delta_{it}b_sz_j+\delta_{jt}b_sz_i+\delta_{is}b_tz_j+\delta_{js}b_tz_i\Big)\mathscr{P}(Z,Z')\\&=\frac{1}{2\pi}\Big(-3\delta_{js}\delta_{it}-3\delta_{jt}\delta_{is}+\frac{1}{2}\delta_{it}z_jb_s+\frac{1}{2}\delta_{jt}z_ib_s+\frac{1}{2}\delta_{is}z_jb_t+\frac{1}{2}\delta_{js}z_ib_t+\frac{1}{4}z_iz_jb_sb_t\Big)\mathscr{P}(Z,Z').\end{aligned} \tag{4.46}$$

By (1.7), (4.45) and (4.46), we get

$$\begin{aligned}
&\Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}z_iz_j\overline{z}_s\overline{z}_t\mathscr{P}\Big)(Z,Z') = \Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}z_iz_j\big(\frac{b_s}{2\pi}+\overline{z}'_s\big)\big(\frac{b_t}{2\pi}+\overline{z}'_t\big)\mathscr{P}\Big)(Z,Z')\\
&= \frac{1}{4\pi^2}\Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}z_iz_jb_sb_t\mathscr{P}\Big)(Z,Z') + \frac{1}{2\pi}\Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}\big(z_iz_j\overline{z}'_tb_s + z_iz_j\overline{z}'_sb_t\big)\mathscr{P}\Big)(Z,Z')\\
&= \frac{1}{4\pi^2}\Big\{-\frac{3}{2\pi}\delta_{it}\delta_{js} - \frac{3}{2\pi}\delta_{jt}\delta_{is} + \frac{1}{2}\delta_{it}z_j\big(\overline{z}_s-\overline{z}'_s\big) + \frac{1}{2}\delta_{jt}z_i\big(\overline{z}_s-\overline{z}'_s\big) + \frac{1}{2}\delta_{is}z_j\big(\overline{z}_t-\overline{z}'_t\big)\\
&\qquad + \frac{1}{2}\delta_{js}z_i\big(\overline{z}_t-\overline{z}'_t\big) + \frac{\pi}{2}z_iz_j\big(\overline{z}_t-\overline{z}'_t\big)\big(\overline{z}_s-\overline{z}'_s\big)\Big\}\mathscr{P}(Z,Z')\\
&\quad + \frac{1}{8\pi^2}\Big\{\big[-2\delta_{is}z_j - 2\delta_{js}z_i + 2\pi z_iz_j(\overline{z}_s-\overline{z}'_s)\big]\overline{z}'_t\\
&\qquad + \big[-2\delta_{it}z_j - 2\delta_{jt}z_i + 2\pi z_iz_j(\overline{z}_t-\overline{z}'_t)\big]\overline{z}'_s\Big\}\mathscr{P}(Z,Z').
\end{aligned}\tag{4.47}$$

By (3.11), (4.44) and (4.47), we get

$$\begin{aligned}
9(d_xd_yI_{26})(0,0) = &-\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_i}}J\big)\frac{\partial}{\partial z_j},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_s}}J\big)\frac{\partial}{\partial\overline{z}_t}\Big\rangle\Big[3(\delta_{it}\delta_{js}+\delta_{jt}\delta_{is})(-2\sqrt{-1})\omega(x_0)\\
&+\delta_{it}dz_j\wedge d\overline{z}_s + \delta_{jt}dz_i\wedge d\overline{z}_s + \delta_{is}dz_j\wedge d\overline{z}_t + \delta_{js}dz_i\wedge d\overline{z}_t\\
&+2\delta_{is}dz_j\wedge d\overline{z}_t + 2\delta_{js}dz_i\wedge d\overline{z}_t + 2\delta_{it}dz_j\wedge d\overline{z}_s + 2\delta_{jt}dz_i\wedge d\overline{z}_s\Big].
\end{aligned}\tag{4.48}$$

By (4.48),

$$\begin{aligned}
(d_xd_yI_{26})(0,0) = &\frac{2}{3}\sqrt{-1}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_i}}J\big)\frac{\partial}{\partial z_j},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\frac{\partial}{\partial\overline{z}_j}+\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_j}}J\big)\frac{\partial}{\partial\overline{z}_i}\Big\rangle\omega(x_0)\\
&-\frac{1}{3}\Big[\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_i}}J\big)\frac{\partial}{\partial z_r},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\frac{\partial}{\partial\overline{z}_q}+\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_q}}J\big)\frac{\partial}{\partial\overline{z}_i}\Big\rangle\\
&\qquad+\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_r}}J\big)\frac{\partial}{\partial z_i},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\frac{\partial}{\partial\overline{z}_q}+\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_q}}J\big)\frac{\partial}{\partial\overline{z}_i}\Big\rangle\Big]dz_r\wedge d\overline{z}_q.
\end{aligned}\tag{4.49}$$

Note that for $U,V,W\in TX$, $\big\langle JU,V\big\rangle=\omega(U,V)$, thus (cf. [13, (8.3.48)]),

$$\big\langle(\nabla^X_UJ)V,W\big\rangle+\big\langle(\nabla^X_VJ)W,U\big\rangle+\big\langle(\nabla^X_WJ)U,V\big\rangle = d\omega(U,V,W)=0.\tag{4.50}$$

From Lemma 3.1, (4.50) and $\big|\frac{\partial}{\partial z_j}\big|^2=\frac{1}{2}$, we have

$$\begin{aligned}
&\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_i}}J\big)\frac{\partial}{\partial z_r},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_q}}J\big)\frac{\partial}{\partial\overline{z}_i}\Big\rangle = 2\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_i}}J\big)\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_r}\Big\rangle\Big[\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\frac{\partial}{\partial\overline{z}_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle-\Big\langle\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_j}}J\big)\frac{\partial}{\partial\overline{z}_i},\frac{\partial}{\partial\overline{z}_q}\Big\rangle\Big].
\end{aligned}\tag{4.51}$$

When we sum (4.51) over $r=q$, we get by (4.42) (cf. [13, (8.3.58)]),

$$\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_i}}J\big)\frac{\partial}{\partial z_q},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_q}}J\big)\frac{\partial}{\partial\overline{z}_i}\Big\rangle=\frac{1}{16}\big|\nabla^XJ\big|^2.\tag{4.52}$$

By (4.7) and (4.51), we get

$$\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_i}}J\big)\frac{\partial}{\partial z_r},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\frac{\partial}{\partial\overline{z}_q}+\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_q}}J\big)\frac{\partial}{\partial\overline{z}_i}\Big\rangle=2\mathscr{J}_{ijr}\big(2\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}-\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}\big).\tag{4.53}$$

By Lemma 3.1, (4.50) and $\big|\frac{\partial}{\partial z_j}\big|^2=\frac{1}{2}$, we obtain

(4.54)
$$\begin{aligned}
&\Big\langle (\nabla^X_{\frac{\partial}{\partial z_r}}J)\frac{\partial}{\partial z_i}, (\nabla^X_{\frac{\partial}{\partial \overline{z}_q}}J)\frac{\partial}{\partial \overline{z}_i}\Big\rangle\\
&= 2\Big[\Big\langle (\nabla^X_{\frac{\partial}{\partial z_i}}J)\frac{\partial}{\partial z_j}, \frac{\partial}{\partial z_r}\Big\rangle + \Big\langle (\nabla^X_{\frac{\partial}{\partial z_j}}J)\frac{\partial}{\partial z_r}, \frac{\partial}{\partial z_i}\Big\rangle\Big]\Big[\Big\langle (\nabla^X_{\frac{\partial}{\partial \overline{z}_i}}J)\frac{\partial}{\partial \overline{z}_j}, \frac{\partial}{\partial \overline{z}_q}\Big\rangle + \Big\langle (\nabla^X_{\frac{\partial}{\partial \overline{z}_j}}J)\frac{\partial}{\partial \overline{z}_q}, \frac{\partial}{\partial \overline{z}_i}\Big\rangle\Big]\\
&= 4\mathscr{J}_{ijr}\big(\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}} - \mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}\big).
\end{aligned}$$

By taking the conjugation of (4.51), we get

$$\Big\langle (\nabla^X_{\frac{\partial}{\partial z_r}}J)\frac{\partial}{\partial z_i}, (\nabla^X_{\frac{\partial}{\partial \overline{z}_i}}J)\frac{\partial}{\partial \overline{z}_q}\Big\rangle = 2\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}\big(\mathscr{J}_{ijr} - \mathscr{J}_{jir}\big) = 2\mathscr{J}_{ijr}\big(\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}} - \mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}\big). \tag{4.55}$$

Substituting (4.42), (4.52), (4.53), (4.54) and (4.55) into (4.49) yields

$$(d_xd_yI_{26})(0,0) = \frac{\sqrt{-1}}{8}\big|\nabla^X J\big|^2\omega(x_0) - \frac{2}{3}\mathscr{J}_{ijr}\big(5\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}} - 4\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}\big)dz_r\wedge d\overline{z}_q. \tag{4.56}$$

4.3. **Evaluation of $(d_xd_yI_2)(0,0)$: part II.** We evaluate now the contribution of I_{22}, I_{24} in $(d_xd_yI_2)(0,0)$. The definitions of $\nabla^X\nabla^X J$ and R^{TX} imply that for $U,V,W,Y\in TX$ (cf. [13, (8.3.59)]),

$$\begin{aligned}
&\big(\nabla^X\nabla^X J\big)_{(U,V)} - \big(\nabla^X\nabla^X J\big)_{(V,U)} = \big[R^{TX}(U,V), J\big],\\
&\Big\langle \big(\nabla^X\nabla^X J\big)_{(Y,U)}V, W\Big\rangle + \Big\langle \big(\nabla^X\nabla^X J\big)_{(Y,V)}W, U\Big\rangle + \Big\langle \big(\nabla^X\nabla^X J\big)_{(Y,W)}U, V\Big\rangle = 0.
\end{aligned} \tag{4.57}$$

Recall that [13, (8.3.71)],

$$\sum_{|\alpha|=2}\big(\partial^\alpha R^L\big)(\mathcal{R}, \frac{\partial}{\partial \overline{z}_i})\frac{Z^\alpha}{\alpha!} = -\sqrt{-1}\pi\Big\langle \big(\nabla^X\nabla^X J\big)_{(\mathcal{R},\mathcal{R})}\mathcal{R}, \frac{\partial}{\partial \overline{z}_i}\Big\rangle - \frac{2\pi}{3}\Big\langle R^{TX}(z,\overline{z})\mathcal{R}, \frac{\partial}{\partial \overline{z}_i}\Big\rangle. \tag{4.58}$$

By (4.50) and (4.57), we get (cf.[13, (8.3.61)]): for $u_1,u_2,u_3\in T^{(1,0)}X$, $\overline{v}_1,\overline{v}_2\in T^{(0,1)}X$,

$$\begin{aligned}
&(\nabla^X\nabla^X J)_{(u_1,u_2)}u_3,\ (\nabla^X\nabla^X J)_{(\overline{v}_1,\overline{v}_2)}u_3 \in T^{(0,1)}X,\\
&(\nabla^X\nabla^X J)_{(u_1,\overline{v}_2)}u_3 \in T^{(1,0)}X,\\
&2\sqrt{-1}\left\langle (\nabla^X\nabla^X J)_{(u_1,\overline{v}_1)}u_2, \overline{v}_2\right\rangle = \left\langle (\nabla^X_{u_1}J)u_2, (\nabla^X_{\overline{v}_1}J)\overline{v}_2\right\rangle .
\end{aligned} \tag{4.59}$$

In particular, we have

$$\begin{aligned}
&\Big\langle \big(\nabla^X\nabla^X J\big)_{(\overline{z},z)}\overline{z}, \frac{\partial}{\partial \overline{z}_i}\Big\rangle = 0,\\
&\Big\langle \big(\nabla^X\nabla^X J\big)_{(z,z)}u_1, \frac{\partial}{\partial \overline{z}_i}\Big\rangle = \Big\langle \big(\nabla^X\nabla^X J\big)_{(\overline{z},\overline{z})}u_1, \frac{\partial}{\partial \overline{z}_i}\Big\rangle = 0,\\
&\Big\langle \big(\nabla^X\nabla^X J\big)_{(\overline{z},z)}u_1, \frac{\partial}{\partial \overline{z}_i}\Big\rangle = \Big\langle \big(\nabla^X\nabla^X J\big)_{(z,\overline{z})}u_1 - [R^{TX}(z,\overline{z}), J]u_1, \frac{\partial}{\partial \overline{z}_i}\Big\rangle\\
&\qquad\qquad = \Big\langle \big(\nabla^X\nabla^X J\big)_{(z,\overline{z})}u_1, \frac{\partial}{\partial \overline{z}_i}\Big\rangle.
\end{aligned} \tag{4.60}$$

By (4.57) and (4.60), we get

$$\begin{aligned}\Big\langle \big(\nabla^X\nabla^X J\big)_{(z,\overline{z})}\overline{z}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle =& \Big\langle \big[R^{TX}(z,\overline{z}), J\big]\overline{z}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle \\ =& -2\sqrt{-1}\Big\langle R^{TX}(z,\overline{z})\overline{z}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle.\end{aligned} \tag{4.61}$$

By (4.57) and (4.59) we get (cf. [13, (8.3.62)]),

$$\Big\langle \big(\nabla^X\nabla^X J\big)_{(u_1,u_2)}\overline{z}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle = \frac{1}{2\sqrt{-1}}\Big\langle \big(\nabla^X_{u_1}J\big)u_2, \big(\nabla^X_{\overline{z}}J\big)\frac{\partial}{\partial\overline{z}_i} - \big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\overline{z}\Big\rangle. \tag{4.62}$$

By (4.59), (4.60), (4.61) and (4.62), we get

$$\begin{aligned}-\pi\sqrt{-1}\Big\langle \big(\nabla^X\nabla^X J\big)_{(\mathcal{R},\mathcal{R})}\mathcal{R}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle =& -\frac{\pi}{2}\Big\langle \big(\nabla^X_z J\big)z, 3\big(\nabla^X_{\overline{z}}J\big)\frac{\partial}{\partial\overline{z}_i} - \big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\overline{z}\Big\rangle \\ &- 2\pi\Big\langle R^{TX}(z,\overline{z})\overline{z}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle - \pi\sqrt{-1}\Big\langle \big(\nabla^X\nabla^X J\big)_{(\overline{z},\overline{z})}\overline{z}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle.\end{aligned} \tag{4.63}$$

By (4.21), (4.58) and (4.63), we get

$$\begin{aligned}I_{22}(Z,Z') =& -\Big[\frac{\pi}{4}\Big\langle \big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_k}, 3\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_s}}J\big)\frac{\partial}{\partial\overline{z}_i} - \big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\frac{\partial}{\partial\overline{z}_s}\Big\rangle \\ &+ \frac{\pi}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j}, \frac{\partial}{\partial\overline{z}_s}\Big)\frac{\partial}{\partial z_k}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle\Big]\Big(\mathscr{L}^{-1}b_i z_j z_k\overline{z}_s\mathscr{P}\Big)(Z,Z') \\ &- \frac{4\pi}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j}, \frac{\partial}{\partial\overline{z}_s}\Big)\frac{\partial}{\partial\overline{z}_t}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle\Big(\mathscr{L}^{-1}b_i z_j\overline{z}_s\overline{z}_t\mathscr{P}\Big)(Z,Z') \\ &- \frac{\pi}{2}\sqrt{-1}\Big\{\mathscr{L}^{-1}b_i\Big\langle \big(\nabla^X\nabla^X J\big)_{(\overline{z},\overline{z})}\overline{z}, \frac{\partial}{\partial\overline{z}_i}\Big\rangle\mathscr{P}\Big\}(Z,Z').\end{aligned} \tag{4.64}$$

By (1.7) and (1.8), we get

$$\begin{aligned}b_i z_j z_k\overline{z}_s\mathscr{P} =& b_i z_j z_k\big(\frac{b_s}{2\pi} + \overline{z}'_s\big)\mathscr{P} \\ =& \Big(\frac{b_i b_s}{2\pi}z_j z_k + \frac{\delta_{js}}{\pi}b_i z_k + \frac{\delta_{ks}}{\pi}b_i z_j + b_i z_j z_k\overline{z}'_s\Big)\mathscr{P}.\end{aligned} \tag{4.65}$$

Thus, by Theorem 1.1, (1.7), (1.8) and (4.65), as in (4.47), we get

$$\begin{aligned}&\Big(\mathscr{L}^{-1}b_i z_j z_k\overline{z}_s\mathscr{P}\Big)(Z,Z') \\ &= \Big(\frac{b_i b_s}{16\pi^2}z_j z_k + \frac{\delta_{js}}{4\pi^2}b_i z_k + \frac{\delta_{ks}}{4\pi^2}b_i z_j + \frac{b_i}{4\pi}z_j z_k\overline{z}'_s\Big)\mathscr{P}(Z,Z') \\ &= \frac{1}{4\pi^2}\Big(-\delta_{js}\delta_{ik} + \pi\delta_{js}z_k\big(\overline{z}_i - \overline{z}'_i\big) - \delta_{ks}\delta_{ij} + \pi\delta_{ks}z_j\big(\overline{z}_i - \overline{z}'_i\big) \\ &\qquad - \pi\delta_{ij}z_k\big(\overline{z}_s - \overline{z}'_s\big) - \pi\delta_{ik}z_j\big(\overline{z}_s - \overline{z}'_s\big) + \pi^2 z_j z_k\big(\overline{z}_i - \overline{z}'_i\big)\big(\overline{z}_s - \overline{z}'_s\big)\Big)\mathscr{P}(Z,Z') \\ &\quad + \frac{1}{4\pi}\Big(-2\delta_{ij}z_k - 2\delta_{ik}z_j + 2\pi z_j z_k\big(\overline{z}_i - \overline{z}'_i\big)\Big)\overline{z}'_s\mathscr{P}(Z,Z').\end{aligned} \tag{4.66}$$

By (4.6) and (4.66), we get

$$\Big(\mathscr{L}^{-1}b_iz_jz_k\overline{z}_s\mathscr{P}\Big)(Z,Z')\approx-\Big[\frac{1}{4\pi^2}\big(\delta_{js}\delta_{ik}+\delta_{ks}\delta_{ij}\big)+\frac{1}{4\pi}\big(\delta_{js}z_k\overline{z}'_i+\delta_{ks}z_j\overline{z}'_i+\delta_{ij}z_k\overline{z}'_s+\delta_{ik}z_j\overline{z}'_s\big)\Big]\mathscr{P}(Z,Z'). \tag{4.67}$$

Again by (1.7) and (1.8),

$$\begin{aligned}b_iz_j\overline{z}_s\overline{z}_t\mathscr{P}&=b_iz_j\Big(\frac{b_s}{2\pi}+\overline{z}'_s\Big)\Big(\frac{b_t}{2\pi}+\overline{z}'_t\Big)\mathscr{P}\\&=\Big[\frac{b_ib_sb_t}{4\pi^2}z_j+\frac{1}{2\pi^2}\big(\delta_{js}b_ib_t+\delta_{jt}b_ib_s\big)+\frac{b_i}{2\pi}\big(b_sz_j+2\delta_{js}\big)\overline{z}'_t+\frac{b_i}{2\pi}\big(b_tz_j+2\delta_{jt}\big)\overline{z}'_s+b_iz_j\overline{z}'_s\overline{z}'_t\Big]\mathscr{P}.\end{aligned} \tag{4.68}$$

By Theorem 1.1 and (4.68), we get

$$\mathscr{L}^{-1}b_iz_j\overline{z}_s\overline{z}_t\mathscr{P}=\Big[\frac{b_ib_sb_t}{48\pi^3}z_j+\frac{1}{16\pi^3}\big(\delta_{js}b_ib_t+\delta_{jt}b_ib_s\big)+\frac{b_i}{16\pi^2}\big(b_sz_j+4\delta_{js}\big)\overline{z}'_t+\frac{b_i}{16\pi^2}\big(b_tz_j+4\delta_{jt}\big)\overline{z}'_s+\frac{b_i}{4\pi}z_j\overline{z}'_s\overline{z}'_t\Big]\mathscr{P}. \tag{4.69}$$

By (1.7), (1.8) and (4.6), we get

$$\Big(\frac{b_i}{16\pi^2}\big(b_sz_j+4\delta_{js}\big)\overline{z}'_t\mathscr{P}\Big)(Z,Z')\approx\frac{1}{4\pi}\big(\delta_{js}\overline{z}_i-\delta_{ij}\overline{z}_s\big)\overline{z}'_t\mathscr{P}(Z,Z'). \tag{4.70}$$

By (4.32), (4.69) and (4.70), we obtain

$$\begin{aligned}\Big(d_xd_y\big(\mathscr{L}^{-1}b_iz_j\overline{z}_s\overline{z}_t\mathscr{P}\big)\Big)(0,0)&=\frac{1}{4\pi}\big(\delta_{js}d\overline{z}_i-\delta_{ij}d\overline{z}_s\big)\wedge d\overline{z}_t+\frac{1}{4\pi}\big(\delta_{jt}d\overline{z}_i-\delta_{ij}d\overline{z}_t\big)\wedge d\overline{z}_s\\&=\frac{1}{4\pi}\big(\delta_{js}d\overline{z}_i\wedge d\overline{z}_t+\delta_{jt}d\overline{z}_i\wedge d\overline{z}_s\big).\end{aligned} \tag{4.71}$$

Finally, by Theorem 1.1, (1.7) and (1.8), as in (4.29), we get

$$\begin{aligned}\Big(\mathscr{L}^{-1}b_i\overline{z}_j\overline{z}_j\overline{z}_t\mathscr{P}\Big)(Z,Z')&=\Big(\mathscr{L}^{-1}b_i\big(\frac{b_j}{2\pi}+\overline{z}'_j\big)\big(\frac{b_s}{2\pi}+\overline{z}'_s\big)\big(\frac{b_t}{2\pi}+\overline{z}'_t\big)\mathscr{P}\Big)(Z,Z')\\&\sim\Big\{\mathscr{L}^{-1}b_i\Big[\frac{b_jb_sb_t}{8\pi^3}+\frac{1}{4\pi^2}\big(b_jb_s\overline{z}'_t+b_jb_t\overline{z}'_s+b_sb_t\overline{z}'_j\big)\Big]\mathscr{P}\Big\}(Z,Z')\\&\approx0.\end{aligned} \tag{4.72}$$

From (3.11), (4.64), (4.67), (4.71) and (4.72), we obtain

$$
\begin{aligned}
(4.73)\quad (d_xd_yI_{22})(0,0)=&\bigg\{\frac{1}{8\pi}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_i},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_j}}J\big)\frac{\partial}{\partial\overline{z}_i}+\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_i}}J\big)\frac{\partial}{\partial\overline{z}_j}\Big\rangle\\
&+\frac{1}{12\pi}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_j}+R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\bigg\}\big(-2\pi\sqrt{-1}\big)\omega(x_0)\\
&+\frac{1}{8}\bigg[\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_r},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_j}}J\big)\frac{\partial}{\partial\overline{z}_q}+\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_q}}J\big)\frac{\partial}{\partial\overline{z}_j}\Big\rangle\\
&+\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_r}}J\big)\frac{\partial}{\partial z_j},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_j}}J\big)\frac{\partial}{\partial\overline{z}_q}+\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_q}}J\big)\frac{\partial}{\partial\overline{z}_j}\Big\rangle\bigg]dz_r\wedge d\overline{z}_q\\
&+\frac{1}{6}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial z_r}+R^{TX}\Big(\frac{\partial}{\partial z_r},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle dz_r\wedge d\overline{z}_q\\
&+\frac{1}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial\overline{z}_r}+R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_r}\Big)\frac{\partial}{\partial\overline{z}_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle d\overline{z}_r\wedge d\overline{z}_q.
\end{aligned}
$$

Note that by (4.23),

$$
(4.74)\qquad R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial z_r}=R^{TX}\Big(\frac{\partial}{\partial z_r},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial z_j}+R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_r}\Big)\frac{\partial}{\partial\overline{z}_j}.
$$

By (4.23), (4.42), (4.52), (4.53), (4.54), (4.55), (4.73) and (4.74), we get

$$
\begin{aligned}
(4.75)\quad (d_xd_yI_{22})(0,0)=&-\sqrt{-1}\bigg[\frac{5}{96}\big|\nabla^XJ\big|^2+\frac{1}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\bigg]\omega(x_0)\\
&+\frac{1}{4}\mathscr{J}_{jir}\big(5\,\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}-4\,\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}\big)dz_r\wedge d\overline{z}_q\\
&+\frac{1}{6}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_r}\Big)\frac{\partial}{\partial\overline{z}_j}+2R^{TX}\Big(\frac{\partial}{\partial z_r},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle dz_r\wedge d\overline{z}_q\\
&+\frac{1}{3}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_j}\Big)\frac{\partial}{\partial\overline{z}_r}+R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_r}\Big)\frac{\partial}{\partial\overline{z}_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle d\overline{z}_r\wedge d\overline{z}_q.
\end{aligned}
$$

By (4.57), (4.59) and (4.60), we get (cf. [14, (2.28)]),

$$
\begin{aligned}
(4.76)\quad \Big\langle\big(\nabla^X\nabla^XJ\big)_{(\mathcal{R},\mathcal{R})}\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big\rangle&=2\Big\langle\big(\nabla^X\nabla^XJ\big)_{(z,\overline{z})}\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_i}\Big\rangle\\
&=-\sqrt{-1}\Big\langle\big(\nabla^X_zJ\big)\frac{\partial}{\partial z_i},\big(\nabla^X_{\overline{z}}J\big)\frac{\partial}{\partial\overline{z}_i}\Big\rangle.
\end{aligned}
$$

By (4.25) and (4.35), we get

$$
(4.77)\qquad \Big(\mathscr{L}^{-1}\mathscr{P}^{\perp}z_s\overline{z}_t\mathscr{P}\Big)(Z,Z')=-\frac{1}{4\pi^2}\Big(\delta_{st}-\pi z_s\big(\overline{z}_t-\overline{z}'_t\big)\Big)\mathscr{P}(Z,Z').
$$

By (3.11), (4.42), (4.54), (4.76) and (4.77), we obtain

$$\begin{aligned}(d_xd_yI_{24})(0,0)&=\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_i},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_j}}J\big)\frac{\partial}{\partial\overline{z}_i}\Big\rangle(-\sqrt{-1})\omega(x_0)\\&\quad+\frac{1}{2}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_s}}J\big)\frac{\partial}{\partial z_i},\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_t}}J\big)\frac{\partial}{\partial\overline{z}_i}\Big\rangle dz_s\wedge d\overline{z}_t\\&=-\frac{\sqrt{-1}}{8}\big|\nabla^XJ\big|^2\omega(x_0)+2\mathscr{J}_{ijr}\big(\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}-\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}\big)dz_r\wedge d\overline{z}_q.\end{aligned}\tag{4.78}$$

Combining (4.22), (4.23), (4.33), (4.38), (4.42), (4.43), (4.56), (4.74), (4.75) and (4.78), we obtain

$$\begin{aligned}-(d_xd_yI_2)(0,0)&=\sqrt{-1}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\omega(x_0)\\&\quad-\frac{1}{2}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_r}\Big)\frac{\partial}{\partial\overline{z}_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle dz_r\wedge d\overline{z}_q-\frac{1}{12}\mathscr{J}_{jir}\big(\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}+4\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}\big)dz_r\wedge d\overline{z}_q.\end{aligned}\tag{4.79}$$

By Lemma 3.1, (4.50), (4.57), (4.51), (4.54), (4.55) and (4.62), we get (cf. [13, (8.3.63)])

$$\begin{aligned}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_r}\Big)\frac{\partial}{\partial\overline{z}_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle&=\frac{\sqrt{-1}}{2}\Big\langle\Big[R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_r}\Big),J\Big]\frac{\partial}{\partial\overline{z}_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle\\&=\frac{\sqrt{-1}}{2}\Big\langle\Big[\big(\nabla^X\nabla^XJ\big)_{(\frac{\partial}{\partial z_j},\frac{\partial}{\partial z_r})}-\big(\nabla^X\nabla^XJ\big)_{(\frac{\partial}{\partial z_r},\frac{\partial}{\partial z_j})}\Big]\frac{\partial}{\partial\overline{z}_j},\frac{\partial}{\partial\overline{z}_q}\Big\rangle\\&=\frac{1}{4}\Big\langle\big(\nabla^X_{\frac{\partial}{\partial z_j}}J\big)\frac{\partial}{\partial z_r}-\big(\nabla^X_{\frac{\partial}{\partial z_r}}J\big)\frac{\partial}{\partial z_j},\ \big(\nabla^X_{\frac{\partial}{\partial\overline{z}_j}}J\big)\frac{\partial}{\partial\overline{z}_q}-\big(\nabla^X_{\frac{\partial}{\partial\overline{z}_q}}J\big)\frac{\partial}{\partial\overline{z}_j}\Big\rangle\\&=\frac{1}{2}\mathscr{J}_{jri}\mathscr{J}_{\overline{j}\overline{q}\overline{i}}.\end{aligned}\tag{4.80}$$

Substituting (4.80) into (4.79) yields

$$(d_xd_yI_2)(0,0)=-\sqrt{-1}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\omega(x_0)+\frac{1}{3}\mathscr{J}_{jir}\big(\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}+\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}\big)dz_r\wedge d\overline{z}_q.\tag{4.81}$$

4.4. **Proof of Theorem 3.3.** By (4.4) and (4.81), we get

$$(d_xd_yI_4)(0,0)=(d_xd_yI_2)(0,0).\tag{4.82}$$

Substituting (4.12), (4.13), (4.17), (4.19), (4.81) and (4.82) into (4.2), we finally obtain

$$\begin{aligned}\big(d_xd_y\mathscr{F}_2\big)(0,0)&=-\frac{6}{9}\mathscr{J}_{jir}\big(\mathscr{J}_{\overline{i}\,\overline{j}\overline{q}}+\mathscr{J}_{\overline{j}\,\overline{i}\overline{q}}\big)dz_r\wedge d\overline{z}_q+2(d_xd_yI_2)(0,0)\\&=-2\sqrt{-1}\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle\omega(x_0).\end{aligned}\tag{4.83}$$

By [13, Theorem 8.3.4, Lemma 8.3.10],

$$8\Big\langle R^{TX}\Big(\frac{\partial}{\partial z_j},\frac{\partial}{\partial\overline{z}_i}\Big)\frac{\partial}{\partial z_i},\frac{\partial}{\partial\overline{z}_j}\Big\rangle=r^X+\frac{1}{4}\big|\nabla^XJ\big|^2=8\pi\boldsymbol{b}_1(x_0).\tag{4.84}$$

The identities (4.83) and (4.84) yield Theorem 3.3. This concludes the proof of Theorem 0.1.

REFERENCES

[1] J.-M. Bismut and E. Vasserot, *The asymptotics of the Ray-Singer analytic torsion associated with high powers of a positive line bundle*, Comm. Math. Phys. **125** (1989), 355–367.

[2] D. Borthwick and A. Uribe, *Nearly Kählerian Embeddings of Symplectic Manifolds*, Asian J. Math. **4** (2000), no. 3, 599–620.

[3] L. Boutet de Monvel and V. Guillemin, *The spectral theory of Toeplitz operators*, Ann. of Math. Stud., vol 99, Princeton Univ. Press, Princeton, NJ, 1981.

[4] L. Boutet de Monvel and J. Sjöstrand, *Sur la singularité des noyaux de Bergman et de Szegő*, Journées: Équations aux Dérivées Partielles de Rennes (1975), pp. 123–164. Astérisque, No. 34–35, Soc. Math. France, Paris, 1976.

[5] D. Catlin, *The Bergman kernel and a theorem of Tian*, Analysis and Geometry in Several Complex Variables (Katata, 1997), 1–23, Trends Math., Birkhäuser Boston, Boston, MA, 1999.

[6] X. Dai, K. Liu, and X. Ma, *On the asymptotic expansion of Bergman kernel*, J. Differential Geom. **72** (2006), no. 1, 1–41.

[7] S. K. Donaldson, *Scalar curvature and projective embeddings. I*, J. Differential Geom. **59** (2001), 479–522.

[8] S. K. Donaldson, *Lower bounds on the Calabi functional*, J. Differential Geom. **70** (2005), 453–472.

[9] V. Guillemin and A. Uribe, *The Laplace operator on the n–th tensor power of a line bundle: eigenvalues which are bounded uniformly in n*, Asymptotic Anal. **1** (1988), 105–113.

[10] M. Lejmi and J. Keller, *On the lower bounds of the L^2-norm of the Hermitian scalar curvature*, Preprint, 2017.

[11] X. Ma, *Geometric quantization on Kähler and symplectic manifolds*, in: Proceedings of the International Congress of Mathematicians, vol. II, pp. 785-810, Hindustan Book Agency, New Delhi, 2010.

[12] X. Ma and G. Marinescu, *The Spin^c Dirac operator on high tensor powers of a line bundle*, Math. Z. **240** (2002), no. 3, 651–664.

[13] X. Ma and G. Marinescu, Holomorphic Morse Inequalities and Bergman Kernels, Progress in Mathematics, vol. 254, Birkhäuser Boston, Inc., Boston, MA, 2007.

[14] X. Ma and G. Marinescu, *Generalized Bergman kernels on symplectic manifolds*, Adv. in Math. **217** (2008), no. 4, 1756–1815.

[15] W.-D. Ruan, *Canonical coordinates and Bergman metrics*, Comm. Anal. Geom. **6** (1998), no. 3, 589–631.

[16] B. Shiffman and S. Zelditch, *Asymptotics of almost holomorphic sections of ample line bundles on symplectic manifolds*, J. Reine Angew. Math. **544** (2002), 181–222.

[17] G. Tian, *On a set of polarized Kähler metrics on algebraic manifolds*, J. Differential Geom. **32** (1998), 99–130.

[18] S.-T. Yau, *Nonlinear analysis in geometry*, Enseign. Math. (2) **33** (1987), no. 1-2, 109–158.

[19] S. Zelditch, *Szegő kernels and a theorem of Tian*, Int. Math. Res. Not. **6** (1998), 317–331.

SCHOOL OF MATHEMATICS AND STATISTICS, HUAZHONG UNIVERSITY OF SCIENCE AND TECHNOLOGY, WUHAN, 430074, HUBEI PROVINCE, CHINA

E-mail address: wlu@hust.edu.cn

IMJ-PRG, UNIVERSITÉ PARIS DIDEROT - PARIS 7, UFR DE MATHÉMATIQUES, CASE 7012, 75205 PARIS CEDEX 13, FRANCE

E-mail address: xiaonan.ma@imj-prg.fr

UNIVERISITÄT ZU KÖLN, MATHEMATISCHES INSTITUT, WEYERTAL 86-90, 50931 KÖLN, GERMANY &INSTITUTE OF MATHEMATICS 'SIMION STOILOW', ROMANIAN ACADEMY, BUCHAREST, ROMANIA

E-mail address: gmarines@math.uni-koeln.de